

## **PRIOR AUTHORIZATION CRITERIA**

This list is current as of October 1, 2025, and pertains to the following formularies:

2026 Pharmacy Benefit Dimensions PDP provided by Labor-Management Healthcare Fund Formulary
--

Pharmacy Benefit Dimensions requires you (or your physician) to get prior authorization for certain drugs listed on the formularies above. This means that you will need to get approval from us before you fill your prescriptions. If you do not get approval, we may not cover the drug. These drugs are listed with a “PA” in the Requirements/Notes column on the formularies. This document contains the Prior Authorization requirements that are associated with the formularies listed above.

If you have any questions, please contact our Medicare Member Services Department at 1-800-667-5936 or, for TTY users 711, October 1<sup>st</sup> – March 31<sup>st</sup>: Monday through Sunday from 8 a.m. to 8 p.m. ET, April 1<sup>st</sup> – September 30<sup>th</sup>: Monday through Friday from 8 a.m. to 8 p.m. ET.

Pharmacy Benefit Dimensions is a subsidiary of Independent Health. Independent Health is a PDP with a Medicare contract. Enrollment in Pharmacy Benefit Dimensions PDP depends on contract renewal between Independent Health and CMS.

The formulary may change at any time. You will receive notice when necessary.

## ACTIMMUNE (interferon gamma-1b)

---

### Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ADEMPAS (riociguat)

## Products Affected

- ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5 MG, 2 MG, 2.5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment (Child-Pugh class C), creatinine clearance below 15 mL/min or on dialysis, concurrent use with nitrates or nitric oxide donors in any form, concurrent use with phosphodiesterase inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by right heart catheterization, submission of patient's WHO Group classification, mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), and pregnancy status for female patients of childbearing potential. For pulmonary arterial hypertension (PAH, WHO Group 1), documentation of pulmonary vascular resistance (PVR) greater than 2 Woods units, submission of current or previous therapies used to treat the condition (see Other Criteria). For chronic thromboembolic pulmonary hypertension (CTEPH, WHO Group 4), confirmation of PVR greater than 3 Woods units, evidence of chronic pulmonary embolism on computed tomography or ventilation/perfusion (V/Q) scan, and attestation patient has inoperable disease or has persistent or recurrent disease after CTEPH surgery (pulmonary thromboendarterectomy).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for PAH (WHO Group 1), the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both (1) sildenafil or tadalafil and (2) ambrisentan or bosentan.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ALECENSA (alectinib)

---

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ALUNBRIG (brigatinib)

---

## Products Affected

- ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# AUGTYRO (repotrectinib)

## Products Affected

- AUGTYRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming tumor is ROS1-positive. For other solid tumors, submission of test confirming tumor has a neurotrophic tyrosine receptor kinase (NTRK) gene fusion, attestation patient has progressed following treatment or patient has no satisfactory alternative therapy.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# AUVELITY (dextromethorphan/bupropion)

## Products Affected

- AUVELITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Seizure disorder, current or prior diagnosis of bulimia or anorexia nervosa, severe hepatic impairment, severe renal impairment, administration of monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient has been screened for and does not have bipolar disorder, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to two generic on-formulary antidepressants (e.g., bupropion, SSRI, SNRI).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# AYVAKIT (avapritinib)

## Products Affected

- AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers or strong CYP3A inhibitors. For advanced or indolent systemic mastocytosis, platelet count below $50 \times 10^9/L$ .
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For gastrointestinal stromal tumor (GIST), submission of test result confirming presence of PDGFRA exon 18 mutation. For advanced or indolent systemic mastocytosis, submission of platelet count.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, immunology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# BALVERSA (erdafitinib)

## Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	PD-1/PD-L1 inhibitor-eligible patients who have not received this therapy, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of susceptible FGFR3 genetic alterations, submission of current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	This drug is not recommended for the treatment of patients who are eligible for and have not received prior PD-1 or PD-L1 inhibitor therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# BEMPEDOIC ACID

## Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant pravastatin utilization with doses above 40 mg/day, concomitant simvastatin utilization with doses above 20 mg/day, history of tendon disorders or rupture
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must (1) currently be using a statin (unless contraindicated) plus ezetimibe, or (2) have tried and failed to have an adequate response to or had an intolerance to (a) at least two statins or (b) one statin and ezetimibe. At least one statin previously tried and failed must be a hydrophilic statin. For each annual reauthorization, documentation that the patient remains on previously-used lipid-lowering therapies since the previous approval, unless there is documentation of a new contraindication or intolerance requiring discontinuation of a therapy (or therapies) since the previous approval, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# BENLYSTA (belimumab)

## Products Affected

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe active central nervous system lupus, patients using other biologic medications or intravenous cyclophosphamide
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current therapies used to treat the condition (see Other Criteria). For systemic lupus erythematosus, submission of autoantibody-positive test result for anti-nuclear antibodies (ANA) and/or anti-double-stranded DNA (anti-dsDNA).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to immunology, nephrology, and rheumatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must be using standard therapy, defined as at least one of the following: systemic corticosteroids (e.g., prednisone), antimalarials (e.g., hydroxychloroquine), or immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate mofetil). For each annual reauthorization, confirmation patient is still using some form of standard therapy (as defined above), unless contraindicated, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

## BESREMI (ropeginterferon alfa-2b-njft)

### Products Affected

- BESREMI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History or presence of severe psychiatric disorders (including severe depression or suicidal ideation), history of presence of active serious or untreated autoimmune disease, moderate or severe hepatic impairment (Child-Pugh class B or C), immunosuppressed transplant recipients, severe or unstable cardiovascular disease (e.g., uncontrolled hypertension, NYHA class 2-4 congestive heart failure, serious cardiac arrhythmia, significant coronary artery stenosis, unstable angina), stroke or myocardial infarction within previous 6 months, severe renal impairment (eGFR less than 30 mL/min)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of eGFR, documentation patient has tried and failed to have an adequate response to or had an intolerance/contraindication to hydroxyurea (HU), pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	If patient is taking HU, initially 12 weeks, then 1 year. If patient is not taking HU, 1 year.
<b>Other Criteria</b>	For the first reauthorization in patients using HU at the start of therapy, attestation patient has tapered completely off HU by the end of week 12 is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# BEXAROTENE GEL

---

## Products Affected

- *bexarotene external*

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# BIOLOGIC RESPONSE MODIFIERS

## Products Affected

- KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- OTEZLA ORAL TABLET 30 MG
- OTEZLA ORAL TABLET THERAPY PACK 10 & 20 & 30 MG
- SOTYKTU
- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML SYRINGE
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- TYENNE SUBCUTANEOUS
- *ustekinumab subcutaneous*
- VELSIPITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria). For all drugs managed by this policy except Otezla and Velsipity, submission of baseline latent tuberculosis screening test (Mantoux tuberculin skin test [a.k.a. PPD test] or interferon-gamma release assay [IGRA]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization of a drug managed by this policy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two preferred agents (an adalimumab biosimilar, Cosentyx, Enbrel, Rinvoq, Skyrizi, an ustekinumab biosimilar, and Xeljanz/Xeljanz XR) for the indication submitted, where possible. For all drugs managed by this policy except Otezla and Velsipity, if TB screening test returns a positive result, coverage will be delayed until latent TB is treated. For each annual reauthorization, yearly TB screening test or chest X-ray required for patients who live in, work in, or travel to areas where TB exposure is likely while on treatment or for those who have previously had a positive TB screening test.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# BOSULIF (bosutinib)

## Products Affected

- BOSULIF ORAL CAPSULE 100 MG, 50 MG
- BOSULIF ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For accelerated or blast phase Ph+ CML, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	For initial authorization for accelerated or blast phase Ph+ CML, the patient must have either (1) tried and had an intolerance to dasatinib, imatinib, or nilotinib or (2) resistance to imatinib, defined as (a) failure to achieve or maintain any hematologic improvement within 4 weeks while on imatinib, or (b) failure to achieve a complete hematologic response by 3 months, cytogenetic response by 6 months or major cytogenetic response by 12 months, or (c) progression of disease after a previous cytogenetic or hematologic response, or (d) presence of a genetic mutation in the BCR-ABL gene associated with imatinib resistance.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

## BRAFTOVI/MEKTOVI (encorafenib/binimetinib)

### Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG
- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E (or V600K mutation for a melanoma diagnosis, pregnancy status for female patients of childbearing potential. For metastatic melanoma or metastatic non-small cell lung cancer, confirmation that encorafenib and binimetinib will be co-administered. For metastatic colorectal cancer, confirmation that encorafenib and cetuximab will be co-administered.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# BUTALBITAL-CONTAINING PRODUCTS IN OLDER PATIENTS

---

## Products Affected

- ASCOMP-CODEINE
- *butalbital-acetaminophen oral tablet 50-300 mg, 50-325 mg*
- *butalbital-apap-caff-cod*
- *butalbital-apap-caffeine oral capsule*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*

- TENCON ORAL TABLET 50-325 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation patient has tried and failed a preferred alternative such as ibuprofen or rizatriptan, or has contraindications to all alternatives.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# CABLIVI (caplacizumab-yhdp)

## Products Affected

- CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation drug will be given with plasma exchange and immunosuppressive therapy. If the coverage determination request is not for the patient's first use of caplacizumab, submission of previous aTTP recurrences while on caplacizumab.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology, hematology, and immunology
Coverage Duration	3 months
Other Criteria	If the coverage determination request is not for the patient's first use of caplacizumab, coverage will not be authorized if the patient has had more than 2 recurrences of aTTP while on therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# CALQUENCE (acalabrutinib)

---

## Products Affected

- CALQUENCE ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, coadministration with strong CYP3A inhibitors
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# CAMZYOS (mavacamten)

## Products Affected

- CAMZYOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Left ventricular ejection fraction (LVEF) less than 55%, coadministration with strong CYP2C19 inhibitors, moderate or strong CYP2C19 inducers, moderate or strong CYP3A4 inducers, a non-dihydropyridine (DHP) calcium channel blocker (CCB) plus a beta-blocker, disopyramide, or ranolazine
<b>Required Medical Information</b>	Diagnosis of covered use including all three of the following: (1) attestation patient has exertional symptoms consistent with the definition of NYHA class II or III disease, (2) confirmation of left ventricular (LV) outflow tract obstruction gradient of at least 50 mm Hg either at rest, during Valsalva maneuver testing, or after exercise, and (3) confirmation of LV wall thickness of at least 15 mm or at least 13 mm if condition is familial, submission of current LVEF, current or previous therapies used to treat the condition (see Other Criteria), pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to both a beta-blocker and a non-DHP CCB. For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# CAPRELSA (vandetanib)

## Products Affected

- CAPRELSA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of congenital long QT syndrome, torsades de pointes, uncompensated heart failure, or bradyarrhythmias, QTcF interval greater than 450 msec, hypocalcemia, hypokalemia, hypomagnesemia, coadministration with strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline serum potassium, calcium, magnesium, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), ECG (or QT/QTcF interval), and pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CARGLUMIC ACID

---

## Products Affected

- *carglumic acid oral tablet soluble*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of elevated plasma ammonia level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated plasma ammonia level since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# CERDELGA (eliglustat)

## Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	Ultrarapid CYP2D6 metabolizers, pre-existing cardiac disease, moderate or severe hepatic impairment, long QT syndrome, coadministration with Class Ia or Class III antiarrhythmics. In patients who are poor or intermediate CYP2D6 metabolizers only, mild hepatic impairment.
Required Medical Information	Diagnosis of covered use, submission of CYP2D6 metabolizer status as detected by a test for determining CYP2D6 genotype, liver function testing or Child-Pugh score.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, updated liver function testing or Child-Pugh score since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# CFTR MODULATOR THERAPIES

## Products Affected

- KALYDECO 150 MG, 50-25-37.5 & 75 MG
- ORKAMBI
- SYMDEKO
- TRIKAFTA ORAL TABLET THERAPY PACK 100-50-75 & TRIKAFTA ORAL THERAPY PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers. For Trikafta, severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis of covered use, submission of cystic fibrosis (CF) mutation test confirming presence of CFTR gene mutations as indicated (see Other Criteria).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Initial authorization requires CF mutation test confirming presence of CFTR gene mutations as follows, by drug being requested: (a) for Kalydeco, a mutation predicted to be responsive to ivacaftor based on section 12.1 of the prescribing information, (b) for Orkambi, two copies of the F508del mutation, (c) for Symdeko, two copies of the F508del mutation or at least one mutation predicted to be responsive based on section 12.1 of the prescribing information, (d) for Trikafta, at least one mutation predicted to be responsive based on section 12.1 of the prescribing information or a responsive mutation based on in vitro data.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CHENODAL (chenodiol)

---

## Products Affected

- CHENODAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known hepatocyte dysfunction, bile duct abnormalities such as intrahepatic cholestasis, primary biliary cirrhosis, or sclerosing cholangitis, radiopaque stones, nonvisualizing gallbladder confirmed as nonvisualizing after 2 consecutive single doses of dye, compelling reasons for gallbladder surgery
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to gastroenterology and hepatology
<b>Coverage Duration</b>	24 months
<b>Other Criteria</b>	Safety beyond 24 months is not established and will not be authorized.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

## CHOLBAM (cholic acid)

---

### Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of liver function testing.
Age Restrictions	
Prescriber Restrictions	Restricted to gastroenterology, hepatology, and pediatric gastroenterology
Coverage Duration	Initially 3 months, then 1 year
Other Criteria	For the first reauthorization, documentation of liver function improvement without complete biliary obstruction or persistent clinical or laboratory indications of worsening liver function or cholestasis is required. For each annual reauthorization, updated liver function testing since the previous authorization is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# COMETRIQ (cabozantinib)

## Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline blood pressure reading, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# COPIKTRA (duvelisib)

## Products Affected

- COPIKTRA ORAL CAPSULE 15 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Request as first- or second-line therapy, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of at least two prior systemic therapies tried and failed, submission of pregnancy status for female patients of childbearing potential, attestation patient will receive prophylaxis for <i>Pneumocystis jirovecii</i> pneumonia (PJP) and, if necessary, cytomegalovirus.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# CORTICOTROPIN

## Products Affected

- CORTROPHIN
- CORTROPHIN GEL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Request for IV administration, treatment of patients under 2 years of age in whom congenital infections are suspected, patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, a history of or presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction, or sensitivity to proteins of porcine origin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of blood pressure reading and baseline serum sodium and potassium levels.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 weeks
<b>Other Criteria</b>	For each reauthorization, updated blood pressure, sodium, and potassium levels since the previous authorization are required. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# COTELLIC/ZELBORAF (cobimetinib/vemurafenib)

## Products Affected

- COTELLIC
- ZELBORAF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For cobimetinib, coadministration with moderate or strong CYP3A inhibitors or inducers. For vemurafenib, electrolyte abnormalities that are not correctable, long QT syndrome, coadministration with drugs that prolong the QT interval.
<b>Required Medical Information</b>	Diagnosis of covered use including verification of BRAF V600 mutation as needed for diagnosis, submission of pregnancy status for female patients of childbearing potential. For patients using cobimetinib, submission of left ventricular ejection fraction (LVEF) with a requirement the baseline LVEF is greater than or equal to 50%. For patients using vemurafenib, submission of QTc interval with a requirement the QT interval is less than or equal to 500 msec.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# CRESEMBA (isavuconazonium)

---

## Products Affected

- CRESEMBA ORAL CAPSULE 186 MG

PA Criteria	Criteria Details
Exclusion Criteria	Familial short QT syndrome, coadministration with strong CYP3A inhibitors or inducers
Required Medical Information	Diagnosis of covered use.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, and oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# CYSTEAMINE EYE DROPS

---

## Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation of corneal cysteine crystal deposits as seen on slit-lamp examination.
Age Restrictions	
Prescriber Restrictions	Restricted to metabolic diseases specialty, ophthalmology, and optometry
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# DALFAMPRIDINE

## Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of seizure, moderate or severe renal impairment (CrCl less than or equal to 50 mL/min)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), confirmation that patient is able to walk.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, updated creatinine clearance since the previous authorization and confirmation patient is able to walk is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DASATINIB

## Products Affected

- *dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia, coadministration with proton pump inhibitors or H2 receptor antagonists
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum potassium and magnesium, pregnancy status for female patients of childbearing potential. For adults with resistance or intolerance to prior therapy, documentation of prior therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DAURISMO (glasdegib)

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, confirmation patient will also be receiving cytarabine as part of chemotherapeutic regimen. If patient is under 75 years of age, documentation of comorbidities that preclude use of intensive induction chemotherapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# DEFERASIROX

## Products Affected

- *deferasirox oral tablet soluble*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, estimated glomerular filtration rate (eGFR) less than 40 mL/min, platelet count below $50 \times 10^9/L$ , high-risk myelodysplastic syndromes, advanced malignancies
<b>Required Medical Information</b>	Diagnosis of covered use, submission of complete blood count (CBC), liver function testing (LFT), ferritin, and eGFR from the previous 3 months.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	For each reauthorization, updated ferritin level and platelet count drawn within last 3 months and updated CBC, LFT, and eGFR drawn within the previous 6 months is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DEFERIPRONE

## Products Affected

- *deferiprone*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Absolute neutrophil count (ANC) below $1.5 \times 10^9/L$ , transfusional iron overload in myelodysplastic syndrome or Diamond Blackfan anemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum ferritin levels, ANC, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, updated ferritin level and ANC within last 3 months is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# DIACOMIT (stiripentol)

---

## Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	Requests for monotherapy, moderate or severe renal impairment, moderate or severe hepatic impairment
Required Medical Information	Diagnosis of covered use, confirmation patient is also receiving clobazam.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# DICLOFENAC PATCH

## Products Affected

- *diclofenac epolamine external*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery, use on non-intact or damaged skin resulting from any etiology including exudative dermatitis, eczema, infection lesions, burns, or wounds, pregnancy after 30 weeks gestation
<b>Required Medical Information</b>	Diagnosis of acute pain, defined as short-term pain not lasting longer than a 3-month period.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No



# DRONABINOL

## Products Affected

- *dronabinol*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. If authorization is requested for treatment of nausea and vomiting associated with cancer therapy, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For initial authorization for treatment of nausea and vomiting associated with cancer therapy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least one 5-HT3 receptor antagonist (e.g., granisetron, ondansetron). If the medication is being administered related to cancer treatment and is a full replacement for intravenous administration of antiemetic therapy within 48 hours of cancer treatment, it is covered as a Part B benefit. To be eligible for Part B coverage, the prescribing physician must indicate this on the prescription.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# ENSACOVE (ensartinib)

## Products Affected

- ENSACOVE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Prior ALK-inhibitor use, severe hepatic impairment, coadministration with moderate or strong CYP3A4 inhibitors or inducers or P-glycoprotein inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, attestation patient has not previously received an ALK-inhibitor, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# EOHILIA (budesonide oral suspension)

## Products Affected

- EOHILIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of upper endoscopy with biopsy showing at least 15 eosinophils per high-power field or 60 eosinophils/mm <sup>2</sup> , documentation of signs/symptoms, including but not limited to trouble swallowing, food sticking in esophagus, acid reflux, abdominal or chest pain, or nausea and vomiting, documentation patient has tried and failed at least an 8-week course of proton pump inhibitor therapy (i.e., patient has eosinophilic esophagitis unrelated to gastroesophageal reflux).
Age Restrictions	11 years of age or older
Prescriber Restrictions	Restricted to allergy, gastroenterology, immunology, and otolaryngology/otorhinolaryngology
Coverage Duration	12 weeks
Other Criteria	A maximum of one 12-week course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# EPIDIOLEX (cannabidiol)

---

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## ERIVEDGE (vismodegib)

---

### Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# EVRYSDI (risdiplam)

## Products Affected

- EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use confirmed by genetic testing including either (a) homozygous deletion of SMN1 exon 7 or (b) compound heterozygosity for SMN1 exon 7 deletion and small mutation, documentation of two or more copies of the SMN2 gene by genetic testing.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For first reauthorization, maintenance of or improvement in any motor score or function compared to baseline is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# FINTEPLA (fenfluramine)

---

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	Administration of monoamine oxidase inhibitors within 14 days of initiation
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# GATTEX (teduglutide)

## Products Affected

- GATTEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including confirmation of dependency on parenteral nutrition at least 3 times per week.
Age Restrictions	
Prescriber Restrictions	Restricted to gastroenterology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization for adults 18 years of age or older, submission of reduction in weekly parenteral nutrition/intravenous support volume from baseline and documentation that a colonoscopy or alternate imaging of the entire colon and upper GI endoscopy with polyp removal and showing no active gastrointestinal malignancy is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# GAVRETO (pralsetinib)

## Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inhibitors, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# GEFITINIB

## Products Affected

- *gefitinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 (L858R) substitution mutations, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# GLP-1 AGONISTS

## Products Affected

- MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.5ML, 12.5 MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML, 5 MG/0.5ML, 7.5 MG/0.5ML
- OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/3ML
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS (FORMULATION R2) ORAL TABLET 1.5 MG, 4 MG, 9 MG
- RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG
- TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Off-label use for weight management (see Other Criteria)
Required Medical Information	Diagnosis of type 2 diabetes confirmed through one of the following: (1) medical record, or (2) ICD-10 on medical claims, or (3) laboratory results (verifying a hemoglobin A1c greater than or equal to 6.5%, a fasting plasma glucose greater than or equal to 126 mg/dL, a 2-hour postprandial blood glucose greater than or equal to 200 mg/dL after an oral glucose tolerance test, or a random plasma blood glucose greater than or equal to 200 mg/dL combined with classic signs/symptoms of hyperglycemia or hyperglycemic crisis), attestation patient is not receiving another GLP-1 agonist for the treatment of any condition.
Age Restrictions	Age must be consistent with the prescribing information of the drug and condition being treated
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	These products will not be approved for weight management as this off-label use is currently excluded from coverage under Medicare Part D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## GOMEKLI (mirdametinib)

---

### Products Affected

- GOMEKLI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to neurology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# HEREDITARY ANGIOEDEMA THERAPIES, MAINTENANCE

## Products Affected

- HAEGARDA
- TAKHZYRO SUBCUTANEOUS SOLUTION
- TAKHZYRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 300 MG/2ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for acute hereditary angioedema (HAE) therapy (attacks)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of objective or subjective documentation that prophylactic therapy is medically necessary, including, but not limited to activity of disease and disease burden, the frequency of HAE attacks, and quality of life.
<b>Age Restrictions</b>	Age must be consistent with the prescribing information of the drug
<b>Prescriber Restrictions</b>	Restricted to allergy, dermatology, hematology, and immunology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# HERNEXEOS (zongertinib)

---

## Products Affected

- HERNEXEOS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation the tumor is nonsquamous, submission of test confirming tumor is HER2-positive, pregnancy status for female patients of childbearing potential, documentation of at least one previous systemic therapy that was tried and failed.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# IBRANCE (palbociclib)

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, attestation the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant, pregnancy status for female patients of childbearing potential. For endocrine-resistant breast cancer, submission of genetic tumor testing confirming primary tumor type is PIK3CA-mutated, attestation patient will be taking palbociclib concurrently with inavolisib.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# IBTROZI (taletrectinib)

## Products Affected

- IBTROZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming tumor is ROS1-positive, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to either crizotinib or entrectinib or have contraindications to both crizotinib and entrectinib.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes



# IDHIFA (enasidenib)

---

## Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH2 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# IMBRUVICA (ibrutinib)

## Products Affected

- IMBRUVICA ORAL CAPSULE
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential. For chronic graft-versus-host disease, documentation of treatment failure with any other systemic immunosuppressive agent.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology, oncology, and transplant specialty
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# IMKELDI (imatinib)

---

## Products Affected

- *imkeldi*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of clinical rationale or documentation detailing why the patient cannot use imatinib oral tablets.
Age Restrictions	
Prescriber Restrictions	Restricted to allergy, dermatology, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# INQOVI (decitabine/cedazuridine)

## Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# INREBIC (fedratinib)

## Products Affected

- INREBIC

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, thiamine deficiency, coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of thiamine level (see Other Criteria) and baseline platelet count, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to ruxolitinib. If baseline thiamine level is low, coverage will be delayed until thiamine is repleted.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# INVEGA INJECTABLE (paliperidone injectable suspension)

## Products Affected

- INVEGA HAFYERA
- INVEGA TRINZA INTRAMUSCULAR SUSPENSION  
 PREFILLED SYRINGE 273 MG/0.88ML, 410  
 MG/1.32ML, 546 MG/1.75ML, 819 MG/2.63ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	Diagnosis of covered use. For the 3-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension. For the 6-month injection, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension or at least one 3-month injection of 3-month paliperidone palmitate extended-release injectable suspension.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# ITOVEBI (inavolisib)

## Products Affected

- ITOVEBI ORAL TABLET 3 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming primary tumor type is HR-positive, HER2-negative, and PIK3CA-mutated, submission of current or previous therapies used to treat the condition (see Other Criteria), attestation that patient has locally advanced or metastatic disease, has not experienced disease progression on or following other PI3K inhibitors, and will be taking inavolisib concurrently with fulvestrant and palbociclib, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must have tried and failed to have an adequate response to endocrine therapy (e.g., tamoxifen or an aromatase inhibitor) and must not have received prior chemotherapy for metastatic breast cancer. In addition, documentation of clinical rationale why abemaciclib, palbociclib, or ribociclib combined with endocrine therapy is not suitable for the patient must be submitted.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# JAKAFI (ruxolitinib)

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	For myelofibrosis, platelet count less than $50 \times 10^9/L$
Required Medical Information	Diagnosis of covered use. For myelofibrosis, submission of baseline platelet count. For polycythemia vera, documented intolerance or inadequate response to hydroxyurea. For acute graft-versus-host disease, documented inadequate response to systemic corticosteroids. For chronic graft-versus-host-disease, documented failure of at least one previous line of systemic therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, oncology, and transplant specialty
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# JOENJA (leniolisib)

## Products Affected

- JOENJA

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, moderate or severe hepatic impairment (Child-Pugh class B or C)
Required Medical Information	Diagnosis of covered use including submission of test confirming presence of a pathogenic variant of either PIK3CD or PIK3R1, submission of liver function testing or Child-Pugh score, confirmation of negative pregnancy status for female patients of childbearing potential or attestation from physician patient is not pregnant and will be using a highly effective method of contraception, attestation patient is not currently using an immunosuppressive medication.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit (e.g., normalization of lymphocyte subsets, normalization of lymphadenopathy, reduction in spleen size, etc.) or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# KETOCONAZOLE ORAL

## Products Affected

- *ketoconazole oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute or chronic liver disease, treatment of fungal meningitis or fungal infections of the skin or nails
<b>Required Medical Information</b>	Diagnosis of culture-proven systemic blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis, submission of baseline ALT, AST, total bilirubin, alkaline phosphatase, prothrombin time and INR, prescriber attestation that the potential benefits of therapy outweigh the risks.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# KISQALI (ribociclib)

## Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Congenital long QT syndrome, QTcF interval greater than or equal to 450 msec at treatment initiation, uncorrected hypokalemia or hypomagnesemia, coadministration with strong CYP3A4 inducers or drugs that can prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HR-positive, HER2-negative, submission of QTcF interval, serum potassium and magnesium drawn within the previous 6 months, pregnancy status for female patients of childbearing potential, attestation that the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# KOSELUGO (selumetinib)

## Products Affected

- KOSELUGO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	Initiation: 2-17 years of age. Continuation: 2 years of age or older.
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Selumetinib is indicated in pediatric patients and will not be approved for adults unless the patient started on the medication prior to 18 years of age.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LAZCLUZE (lazertinib)

## Products Affected

- LAZCLUZE ORAL TABLET 240 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations, attestation that the medication will be used in combination with amivantamab and will be given with anticoagulant prophylaxis for the first four months of therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LEUKINE (sargramostim, GM-CSF)

---

## Products Affected

- LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# L-GLUTAMINE

---

## Products Affected

- *l-glutamine oral packet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of sickle cell disease, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	
Prescriber Restrictions	Restricted to hematology
Coverage Duration	1 year
Other Criteria	For initial authorization, the patient must be using, tried and failed to have an adequate response to, or had an intolerance/contraindication to hydroxyurea.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# LIDOCAINE TRANSDERMAL PATCHES

## Products Affected

- *lidocaine external patch 5 %*
- LIDOCAN
- TRIDACAINE II

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# LIVTENCITY (maribavir)

---

## Products Affected

- LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including a documented history of hematopoietic stem cell or solid organ transplant, submission of current or previous therapies used to treat the condition (see Other Criteria).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to hematology, infectious diseases, oncology, and transplant specialty
Coverage Duration	8 weeks
Other Criteria	For authorization, the patient must have tried and failed to have an adequate response to at least one of cidofovir, foscarnet, ganciclovir, or valganciclovir.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LODOCO (colchicine)

## Products Affected

- LODOCO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for the treatment of gout, renal failure, severe hepatic impairment, pre-existing blood dyscrasias, coadministration with strong CYP3A4 or P-glycoprotein inhibitors
<b>Required Medical Information</b>	Diagnosis, documented by either (1) prior acute coronary syndrome, or (2) prior ischemic stroke, transient ischemic attack, or carotid artery stenosis greater than 50%, or (3) prior coronary revascularization, or (4) proven coronary artery disease on invasive coronary angiography or computer tomography angiography, or (5) coronary-artery calcium score greater than or equal to 300 Agatston units, or (6) aortic atherosclerotic disease, or (7) symptomatic peripheral vascular disease, submission of estimated glomerular filtration rate (eGFR) or creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance) with a requirement the eGFR or creatinine clearance is greater than or equal to 15 mL/min, and attestations patient (1) does not have severe hepatic impairment, and (2) has had a recent complete blood count and does not have evidence of any cytopenia, and (3) does not have NYHA functional Class 3 or 4 heart failure.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, submission of updated eGFR or creatinine clearance and complete blood count since the previous authorization is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# LORBRENA (lorlatinib)

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, baseline blood pressure, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# LYBALVI (olanzapine/samidorphan)

## Products Affected

- LYBALVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis, coadministration with opioids, levodopa, dopamine agonists, or strong CYP3A inducers, acute opioid withdrawal, end-stage renal disease
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance to both (1) generic olanzapine, including documentation showing a positive therapeutic benefit but unacceptable weight gain as a result of the drug, and (2) one other generic second-generation antipsychotic with low incidence of metabolic side effects (e.g., aripiprazole, lurasidone, ziprasidone).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# MAVYRET (glecaprevir/pibrentasvir)

## Products Affected

- MAVYRET

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh class B or C), coadministration with rifampin or atazanavir
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Criteria for coverage duration will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# MEGESTROL IN OLDER PATIENTS

## Products Affected

- *megestrol acetate oral suspension 40 mg/ml, 625 mg/5ml*

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	PA not required for hematology or oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# MEKINIST/TAFINLAR (trametinib/dabrafenib)

## Products Affected

- MEKINIST
- TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of BRAF V600E or V600K mutation, pregnancy status for female patients of childbearing potential. For anaplastic thyroid cancer, BRAF V600E-mutated solid tumors, low-grade glioma, and adjuvant BRAF V600E- and/or V600K-mutated melanoma indications, confirmation that trametinib and dabrafenib will be co-administered.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# MEMANTINE/DONEPEZIL

## Products Affected

- *memantine hcl-donepezil hcl*
- NAMZARIC ORAL CAPSULE EXTENDED RELEASE 24 HOUR 7-10 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of documentation that the patient has tolerated donepezil 10 mg daily for a minimum of 1 month.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes



# MODEYSO (dordaviprone)

---

## Products Affected

- MODEYSO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming an H3K27M mutation, pregnancy status for female patients of childbearing potential, documentation of at least one previous therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# NERLYNX (neratinib)

## Products Affected

- NERLYNX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with proton pump inhibitors, strong CYP3A4 inhibitors, moderate CYP3A4 and P-glycoprotein dual inhibitors, or moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HER2-positive, confirmation member has completed adjuvant trastuzumab-based therapy or will be using in combination with capecitabine, pregnancy status for female patients of childbearing potential. For advanced or metastatic breast cancer, submission of previous anti-HER2 regimens used.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NILOTINIB

## Products Affected

- DANZITEN
- *nilotinib hcl*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia, long QT syndrome, coadministration with drugs that prolong the QT interval, proton pump inhibitors, or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation of positive Philadelphia chromosome (Ph) status, baseline serum potassium and magnesium levels.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# NINLARO (ixazomib)

---

## Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, documentation that medication will be administered concomitantly with lenalidomide and dexamethasone, documentation of prior therapy regimen for multiple myeloma, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# NITISINONE

## Products Affected

- *nitisinone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of succinylacetone in urine or plasma.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a clinical benefit, such as reductions in urine succinylacetone level, alpha-fetoprotein level, serum tyrosine level, or serum phenylalanine level, or maintenance of a benefit previously achieved is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# NUPLAZID (pimavanserin)

## Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Psychosis unrelated to Parkinson's disease psychosis, cardiac arrhythmias, symptomatic bradycardia, congenital QT prolongation, coadministration with moderate or strong CYP3A4 inducers or drugs that prolong the QT interval, uncorrected hypokalemia or hypomagnesemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline serum potassium and magnesium levels.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

## ODOMZO (sonidegib)

---

### Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, coadministration with strong CYP3A4 inhibitors or moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation patient is not a candidate for surgery or radiation therapy or carcinoma has recurred following surgery or radiation therapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## OGSIVEO (nirogacestat)

---

### Products Affected

- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inhibitors or inducers, proton pump inhibitors, or H2 receptor antagonists
Required Medical Information	Diagnosis of covered use with documentation of tumor progression, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and sarcoma specialty
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



## OJEMDA (tovorafenib)

### Products Affected

- OJEMDA ORAL SUSPENSION RECONSTITUTED
- OJEMDA ORAL TABLET 100 MG, 100 MG (16 PACK), 100 MG (24 PACK)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP2C8 inhibitors or inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming presence of BRAF V600 mutation or BRAF gene fusion or rearrangement, documentation of previous systemic therapy/therapies for pediatric low-grade glioma tried and failed with a minimum of one previous therapy necessary for approval, pregnancy status for female patients of childbearing potential. If genetic testing does not reveal a BRAF gene fusion or rearrangement, documentation of previous intolerance to, contraindication to, or other reason why the patient cannot use the combination of trametinib and dabrafenib.
<b>Age Restrictions</b>	Initiation: 21 years of age or younger (see Other Criteria)
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Tovorafenib is indicated as therapy in children and young adults and will not be approved for adults unless the patient started on the medication prior to 22 years of age.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

## OJJAARA (mometotinib)

---

### Products Affected

- OJJAARA ORAL TABLET 100 MG, 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active infection, uncontrolled acute or chronic liver disease
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ONUREG (azacitidine)

## Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	Diagnosis of myelodysplastic syndrome
Required Medical Information	Diagnosis of covered use, documentation patient achieved first complete remission or complete remission with incomplete blood count recovery following intensive induction chemotherapy and attestation patient cannot complete intensive curative therapy, submission of absolute neutrophil count (with the requirement it is at least 500/mcL), submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## ORIAHNN (elagolix/estradiol/norethindrone)

### Products Affected

- ORIAHNN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known liver impairment or disease, known osteoporosis, undiagnosed abnormal uterine bleeding, women who are at increased risk of, have a history of, or currently have thrombotic or thromboembolic disorders (including women over 35 years of age who smoke and women with uncontrolled hypertension), current/history of breast cancer or other hormone-sensitive cancer
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is premenopausal, submission of baseline blood pressure, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	2 years (see Other Criteria)
<b>Other Criteria</b>	Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ORILISSA (elagolix)

## Products Affected

- ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment (Child-Pugh class C), known osteoporosis, coadministration with OATP1B1 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is premenopausal and will be using non-hormonal contraception during therapy, submission of liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential. If the patient has previously used elagolix, submission of dose used and number of total months of prior use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	Up to 24 months based on dose and coexisting conditions (see Other Criteria)
<b>Other Criteria</b>	Due to increased risk of bone loss, maximum duration of use is limited based on dose and coexisting conditions. For (1) endometriosis with dyspareunia where dose will be 200 mg twice daily, or (2) women with moderate hepatic impairment, the maximum duration of use is 6 months. Requests for use greater than 6 months will not be approved in these situations. For (3) endometriosis without dyspareunia, 150 mg daily for 24 months. Requests for use greater than 24 months will not be approved in this situation.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# OXERVATE (cenegermin-bkbj)

---

## Products Affected

- OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use confirming Stage 2 or 3 neurotrophic keratitis in at least one eye.
Age Restrictions	
Prescriber Restrictions	Restricted to optometry and ophthalmology
Coverage Duration	8 weeks
Other Criteria	PA applies to all. Safety and efficacy beyond on 8-week course of therapy is not established and will not be authorized.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PANRETIN (alitretinoin)

## Products Affected

- PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy, instances where systemic Kaposi sarcoma therapy is required (more than 10 new Kaposi's sarcoma lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary Kaposi sarcoma, or symptomatic visceral involvement)
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PAZOPANIB

## Products Affected

- *pazopanib hcl*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, uncontrolled hypertension, uncorrected hypokalemia, hypocalcemia, or hypomagnesemia, coadministration with strong CYP3A4 inducers, proton pump inhibitors, H2-receptor antagonists, or drugs that can prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline blood pressure, serum potassium, calcium, and magnesium levels, pregnancy status for female patients of childbearing potential. For soft tissue sarcoma, submission of previous chemotherapy regimen(s).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No



# PEGFILGRASTIM

## Products Affected

- UDENYCA
- UDENYCA ONBODY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of FDA-approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PEMAZYRE (pemigatinib)

---

## Products Affected

- PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of either FGFR1 rearrangement or FGFR2 fusion or rearrangement depending on the indication, attestation patient has used previous systemic treatment for the indication, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PIQRAY (alpelisib)

## Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing confirming the primary tumor type is HR-positive, HER2-negative, and PIK3CA-mutated, attestation that patient has advanced or metastatic disease and will be taking concurrently with fulvestrant, submission of at least one endocrine-based (e.g., anastrozole, exemestane, letrozole, tamoxifen, etc.) regimen tried and failed, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PRETOMANID

## Products Affected

- *pretomanid*

PA Criteria	Criteria Details
Exclusion Criteria	Inability to use bedaquiline or linezolid, drug-sensitive tuberculosis, coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, attestation pretomanid will be used in combination with bedaquiline and linezolid.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to infectious diseases and pulmonology
Coverage Duration	26 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# PREVYMIS (letermovir)

## Products Affected

- PREVYMIS ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), coadministration with ergot alkaloids, pimozide, or pitavastatin or simvastatin when coadministered with cyclosporine
<b>Required Medical Information</b>	Diagnosis of covered use, submission of day number post-transplant. For use after kidney transplant, documentation patient is high risk, defined as donor CMV seropositive/recipient CMV seronegative (D+/R-), submission of explanation why valganciclovir is contraindicated or cannot be used for prophylaxis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology, infectious diseases, oncology, and transplant speciality
<b>Coverage Duration</b>	100 days post-HSCT or 200 days post-kidney transplant or post-HSCT at risk for late CMV infection
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# PROMETHAZINE IN OLDER PATIENTS

## Products Affected

- *promethazine hcl oral solution 6.25 mg/5ml*
- *promethazine hcl oral tablet*
- *promethazine hcl rectal suppository 12.5 mg, 25 mg*
- *promethazine-phenylephrine*
- PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For allergic conditions, documentation must be submitted showing patient has tried and failed or had an inadequate response to a second-generation antihistamine.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Promethazine is a potent anticholinergic considered high-risk in older patients due to risks of confusion, dry mouth, constipation, and decreased clearance with advanced age.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# PROSTATE CANCER ORAL MEDICATIONS

## Products Affected

- AKEEGA
- ERLEADA
- NUBEQA
- XTANDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For Akeega, severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension, uncontrolled hypokalemia
<b>Required Medical Information</b>	Diagnosis of covered use. For Nubeqa, submission of current or previous therapies used to treat the condition (see Other Criteria). For Akeega, submission of test confirming presence of deleterious BRCA mutation, Child-Pugh score or liver function testing, baseline blood pressure reading, and serum potassium level.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology and urology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization of Nubeqa, the patient must have tried and failed to have an adequate response or had an intolerance to both Erleada and Xtandi.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# PULMONARY HYPERTENSION MEDICATIONS

## Products Affected

- ALYQ
- *ambrisentan oral tablet 10 mg, 5 mg*
- *bosentan oral tablet 125 mg, 62.5 mg*
- OPSUMIT
- ORENITRAM
- ORENITRAM MONTH 1
- ORENITRAM MONTH 2
- ORENITRAM MONTH 3
- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*
- *tadalafil (pah)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For ambrisentan, bosentan, or Opsumit, pregnancy. For ambrisentan or Orenitram, moderate or severe hepatic impairment. For tadalafil, severe hepatic impairment or creatinine clearance below 30 mL/min or on hemodialysis. For ambrisentan, idiopathic pulmonary fibrosis.
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by right heart catheterization, submission of mean pulmonary arterial pressure greater than 20 mm Hg at rest, pulmonary arterial wedge pressure less than or equal to 15 mm Hg, pulmonary vascular resistance greater than 2 Woods units, and attestation patient is WHO Group 1. For ambrisentan, bosentan, or Opsumit, submission of pregnancy status for female patients of childbearing potential. For Opsumit only, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	For all drugs in this policy except bosentan, 18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization of Opsumit, the patient must have tried and failed to have an adequate response to or had an intolerance to ambrisentan or bosentan.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes



# RETEVMO (selpercatinib)

## Products Affected

- RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension, coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For RET fusion-positive thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible. For solid tumors with a RET gene fusion, documentation of previous systemic therapy tried or reason why patient has no satisfactory alternative treatment options.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# REVCOVI (elapegademase-lvlr)

## Products Affected

- REVCOVI

PA Criteria	Criteria Details
Exclusion Criteria	Severe thrombocytopenia
Required Medical Information	Diagnosis of covered use confirmed by either biochemical testing showing less than 1% of adenosine deaminase (ADA) catalytic activity in red blood cells or genetic testing showing biallelic ADA pathogenic variants, submission of baseline plasma ADA level (if genetic testing was submitted as confirmation of diagnosis) and platelet count.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, immunology, and specialists in genetic diseases
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, submission of updated plasma ADA level demonstrating an improvement from baseline and platelet count is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# REVUFORJ (revumenib)

## Products Affected

- REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	QTcF interval greater than 450 msec at treatment initiation, white blood cell (WBC) count greater than $25 \times 10^9/L$ , uncorrected hypokalemia or hypomagnesemia, coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test confirming presence of a lysine methyltransferase 2A gene (KMT2A) translocation that is not a 11q23 partial tandem duplication, submission of baseline QTcF interval, serum potassium, serum magnesium, baseline WBC count, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# REZLIDHIA (olutasidenib)

---

## Products Affected

- REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# RIVFLOZA (nedosiran)

## Products Affected

- RIVFLOZA

PA Criteria	Criteria Details
Exclusion Criteria	Estimated glomerular filtration rate (eGFR) less than 30 mL/min/1.73 m <sup>2</sup>
Required Medical Information	Diagnosis of covered use, documentation of AGXT mutation confirmed by liver enzyme analysis or genetic testing, submission of 24-hour urinary oxalate (Uox) excretion with a requirement it is greater than or equal to 0.7 mmol (normalized to body surface area if patient is under 18 years of age) and eGFR, attestation patient has not received a prior kidney or liver transplant, attestation patient will not be using in combination with lumasiran (Oxlumo).
Age Restrictions	
Prescriber Restrictions	Restricted to nephrology and urology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of clinically relevant response to therapy as evidenced by reduced Uox or plasma oxalate levels is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ROMVIMZA (vimseltinib)

## Products Affected

- ROMVIMZA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal
<b>Required Medical Information</b>	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# ROZLYTREK (entrectinib)

## Products Affected

- ROZLYTREK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inducers or drugs that prolong the QTc interval
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of test confirming presence of ROS1-positive tumor. For solid tumors, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation and attestation tumor is metastatic or surgical resection/other systemic therapies are unsatisfactory treatment options.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# RYDAPT (midostaurin)

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For acute myeloid leukemia, submission of test confirming presence of FLT3 mutation, attestation patient will be receiving cytarabine and daunorubicin induction and cytarabine consolidation with midostaurin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to allergy, hematology, and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# SAPROPTERIN

## Products Affected

- JAVYGTOR
- *sapropterin dihydrochloride oral packet*
- *sapropterin dihydrochloride oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of blood phenylalanine concentration.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For the first reauthorization, documentation of reduction in blood phenylalanine concentration from pre-treatment baseline is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## SECOND-GENERATION ANTIPSYCHOTICS

### Products Affected

- CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG, 42 MG
- COBENFY
- COBENFY STARTER PACK
- FANAPT ORAL TABLET 1 MG, 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG
- FANAPT TITRATION PACK
- FANAPT TITRATION PACK A
- FANAPT TITRATION PACK B ORAL TABLET
- FANAPT TITRATION PACK C ORAL TABLET
- REXULTI ORAL TABLET 0.25 MG, 0.5 MG, 1 MG, 2 MG, 3 MG, 4 MG
- SECUADO TRANSDERMAL PATCH 24 HOUR 3.8 MG/24HR, 5.7 MG/24HR, 7.6 MG/24HR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis. For Cobenfy only, moderate to severe renal impairment, urinary retention, gastric retention, hepatic impairment, untreated narrow-angle glaucoma.
<b>Required Medical Information</b>	Diagnosis of covered use. For schizophrenia, an indication related to bipolar disorder type I, or for Rexulti for major depressive disorder, submission of current or previous therapies used to treat the condition (see Other Criteria). For Cobenfy only, submission of estimated glomerular filtration rate with the requirement it is at least 60 mL/min, attestation patient does not have hepatic impairment.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization for schizophrenia or acute treatment of manic/mixed episodes of bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and at least one other generic second-generation atypical antipsychotic. For initial authorization of Caplyta for depressive episodes associated with bipolar I disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two of the following drugs: cariprazine, lurasidone, olanzapine, or quetiapine. For Rexulti as an adjunctive therapy to antidepressants for major depressive disorder, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and cariprazine or quetiapine.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

# SEDATIVE HYPNOTICS IN OLDER PATIENTS

## Products Affected

- *eszopiclone*
- *zaleplon*
- *zolpidem tartrate oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation at least two of the following medications were tried and deemed ineffective or intolerable: Belsomra, doxepin tablets, ramelteon, and trazodone.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	Sedative hypnotic medications are high-risk medications in older patients due to increased risks of cognitive impairment, delirium, unsteady gait, syncope, falls, fractures, and motor vehicle accidents.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# SKYCLARYS (omaveloxolone)

## Products Affected

- SKYCLARYS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by genetic testing, submission of liver function testing or Child-Pugh score.
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and specialists in genetic diseases
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SOFOSBUVIR/VELPATASVIR

---

## Products Affected

- *sofosbuvir-velpatasvir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) infection and HCV genotype, attestation that patients with decompensated cirrhosis will receive concomitant ribavirin therapy unless ribavirin therapy is contraindicated.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# SOMAVERT (pegvisomant)

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use including attestation that surgery or radiation was not curative or is not an option, submission of baseline IGF-1, submission of baseline liver function testing (LFT) including bilirubin with the requirement liver transaminases are less than or equal to 3 times the upper limit of normal (ULN). If liver transaminases are greater than 3 times ULN, submission of the cause of liver dysfunction determined through a comprehensive workup.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For the first reauthorization, submission of updated IGF-1 level demonstrating an improvement from baseline, LFT showing liver transaminases below 5 times the ULN, and attestation patient does not have signs or symptoms of liver injury (e.g., jaundice, elevated bilirubin level or bilirubinuria, fatigue, nausea, vomiting, right upper quadrant pain, ascites, unexplained edema, easy bruisability) is required. For each annual reauthorization, documented improvement/maintenance of IGF-1 level is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# SORAFENIB

## Products Affected

- *sorafenib tosylate*

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For differentiated thyroid cancer, attestation patient is radioactive iodine-refractory or ineligible.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TABRECTA (capmatinib)

---

## Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# TADALAFIL FOR BPH

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Erectile dysfunction indication (see Other Criteria), severe hepatic impairment (Child-Pugh class C), severe renal impairment
<b>Required Medical Information</b>	Diagnosis of benign prostatic hyperplasia (BPH), submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to urology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required. This product will not be approved for erectile dysfunction as this use is currently excluded from coverage under Medicare Part D.
<b>Indications</b>	Some FDA-approved Indications Only.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TAGRISSO (osimertinib)

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions, exon 21 L858R mutations, or T790M mutations, pregnancy status for female patients of childbearing potential. For EGFR T790M mutation-positive NSCLC, submission of previous EGFR tyrosine kinase inhibitor therapy used for indication.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TALZENNA (talazoparib)

## Products Affected

- TALZENNA ORAL CAPSULE 0.1 MG, 0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For breast cancer, submission of test results confirming germline BRCA mutation-positive, human epidermal growth factor receptor 2 (HER2) negative disease. For prostate cancer, submission of test results confirming HRR gene-mutated disease, confirmation talazoparib will be used in combination with enzalutamide.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TASIMELTEON

## Products Affected

- *tasimelteon*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with strong CYP1A2 inhibitors or CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use. For non-24-hour sleep-wake disorder, attestation patient is totally blind. For Smith-Magenis Syndrome, documentation of genetic testing results confirming chromosome 17p11.2 deletion.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to neurology and sleep medicine
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For each annual reauthorization, submission of objective documentation of a symptomatic or clinical benefit or maintenance of a benefit previously achieved is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TAVNEOS (avacopan)

## Products Affected

- TAVNEOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inducers, active serious infection, chronic active hepatitis B, untreated hepatitis C, uncontrolled autoimmune hepatitis, cirrhosis
<b>Required Medical Information</b>	Diagnosis of covered use (granulomatosis with polyangiitis or microscopic polyangiitis variants of anti-neutrophil cytoplasmic antibody [ANCA]-associated vasculitis), submission of HBV serology testing, attestation patient is using rituximab, cyclophosphamide/azathioprine, or another compendium-supported therapy for the treatment of ANCA-associated vasculitis, along with glucocorticoids.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to immunology, nephrology, pulmonology, and rheumatology
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	For the first reauthorization, documentation of clinically relevant response to therapy, including but not limited to disease remission defined using changes in Birmingham Vasculitis Activity Score, a documented reduction in maintenance glucocorticoid dose, or improved or sustained renal function, is required. For each annual reauthorization, documented maintenance of a clinical benefit is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TEPMETKO (tepotinib)

---

## Products Affected

- TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# TERIPARATIDE

## Products Affected

- *teriparatide subcutaneous solution pen-injector 560 mcg/2.24ml, 620 mcg/2.48ml*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pre-existing hypercalcemia, underlying hypercalcemic disorder (such as primary hyperparathyroidism), patients with an increased risk of osteosarcoma (such as those with Paget's disease)
<b>Required Medical Information</b>	Diagnosis of covered use where "high risk for fracture" is defined as documentation of (1) a history of fracture of the hip or vertebra regardless of bone mineral density (BMD), or (2) a history of fracture of the proximal humerus, pelvis, or distal forearm and T-score between -1.0 and -2.5, or (3) T-score less than or equal to -2.5 at the total hip, femoral neck, spine, or distal third of the radius, or (4) T-score between -1.0 and -2.5 at the total hip, femoral neck, spine, or distal third of the radius, and (a) a 10-year probability of hip fracture as assessed by FRAX score of at least 3%, or (b) a 10-year probability of a major osteoporosis-related fracture as assessed by FRAX score of at least 20%, submission of baseline serum calcium, postmenopausal status, current or previous therapies used to treat the condition (see Other Criteria), number of total months of all prior use of parathyroid hormone analogs and parathyroid hormone related peptides.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	2 years unless patient is at high risk for fracture after 2 years of therapy (see Other Criteria)
<b>Other Criteria</b>	For initial authorization, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to at least one bisphosphonate. Therapeutic failure is defined as either a fracture or a decrease in BMD while using a bisphosphonate for at least 3 months. Use of parathyroid hormone analogs and/or parathyroid hormone related peptides for more than 2 years during a patient's lifetime is generally not recommended. For annual reauthorization beyond 2 years, submission of updated serum calcium since initial authorization and evidence the patient remains at high risk for fracture as defined in the Required Medical Information section is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

PA Criteria	Criteria Details
Prerequisite Therapy Required	Yes



# TOLVAPTAN (HYPONATREMIA)

## Products Affected

- *tolvaptan oral tablet 15 mg*tolvaptan (hyponatremia),  
30 mg
tolvaptan (hyponatremia)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Underlying liver disease, need to raise serum sodium acutely, inability to sense or respond to thirst, hypovolemia, anuria, coadministration with strong CYP3A inhibitors or inducers or desmopressin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of evidence of clinically significant hyponatremia, defined as (1) serum sodium less than 125 mEq/L, or (2) serum sodium less than 135 mEq/L that is symptomatic and has resisted correction with fluid restriction.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	30 days (see Other Criteria)
<b>Other Criteria</b>	Treatment should be initiated in a setting where serum sodium can be monitored closely. Treatment is limited to 30 days to prevent liver injury. This formulation of tolvaptan will not be approved for autosomal dominant polycystic kidney disease (ADPKD) as it is not indicated for ADPKD and the tolvaptan formulation approved for ADPKD has a mandatory REMS program, making it only available through a restricted distribution program.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# TURALIO (pexidartinib)

---

## Products Affected

- TURALIO ORAL CAPSULE 125 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal, coadministration with other hepatotoxic medications, strong CYP3A inducers, or proton pump inhibitors
Required Medical Information	Diagnosis of covered use (and documentation surgical intervention is not possible or practical), documentation of patient's severe morbidity or functional limitations, submission of serum transaminases, total and direct bilirubin, and ALP, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VALCHLOR (mechlorethamine)

---

## Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	Use as initial therapy
Required Medical Information	Diagnosis of covered use, submission of previous skin-directed therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VANFLYTA (quizartinib)

## Products Affected

- VANFLYTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for maintenance monotherapy after allogeneic hematopoietic stem cell transplant, uncorrected hypokalemia or hypomagnesemia, long QT syndrome, QTcF interval greater than 450 msec at treatment initiation, coadministration with moderate or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use including submission of test confirming presence of FLT3 internal tandem duplication-positive mutation, submission of QTcF interval, baseline serum potassium and magnesium levels, and pregnancy status for female patients of childbearing potential, attestation patient does not have history of ventricular arrhythmias or torsades de pointes.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	No

# VANRAFIA (atrasentan)

## Products Affected

- VANRAFIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment, coadministration with moderate or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of primary IgA nephropathy confirmed by biopsy, submission of 24-hour urine protein of at least 1 g/day or 24-hour urine protein-to-creatinine ratio (UPCR) of at least 0.8 g/g, liver function testing or Child-Pugh score, pregnancy status for female patients of childbearing potential, submission of current or previous therapies used to treat the condition (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to immunology and nephrology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	For initial authorization, the patient must be currently using (or have a contraindication to the use of) a maximally-tolerated dose of an ACE inhibitor or ARB and tried and failed to have an adequate response to or had an intolerance to at least one immunosuppressant (e.g., azathioprine, mycophenolate, etc.). For each annual reauthorization, documentation of clinically relevant response to therapy, including either stabilization or improvement of UPCR or a reduction in 24-hour urine protein from baseline, is required.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No
<b>Prerequisite Therapy Required</b>	Yes

## VENCLEXTA (venetoclax)

---

### Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers. For CLL/SLL, coadministration with strong CYP3A inhibitors at treatment initiation and initial dosage titration.
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VERZENIO (abemaciclib)

## Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers or ketoconazole
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing confirming that the primary tumor type is HR-positive, HER2-negative, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VIJOICE (alpelisib)

## Products Affected

- VIJOICE ORAL PACKET
- VIJOICE ORAL TABLET THERAPY PACK 125 MG, 200 & 50 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use including at least one target lesion on imaging with requesting provider attestation patient has severe or life-threatening disease, submission of test confirming presence of mutation in PIK3CA gene, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to specialists in genetic diseases or inborn errors of metabolism
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	For each reauthorization, submission of objective documentation of a symptomatic or clinical benefit (e.g., reductions in target lesion size, pain, vascular malformations, limb enlargements, etc.), or maintenance of improvement previously achieved, is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# VITRAKVI (larotrectinib)

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation, attestation tumor is metastatic or surgical resection and other systemic therapies are unsatisfactory treatment options, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VIZIMPRO (dacomitinib)

---

## Products Affected

- VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with a proton pump inhibitor
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of EGFR exon 19 deletions or exon 21 L858R substitution mutations, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VONJO (pacritinib)

## Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh class B or C), estimated glomerular filtration rate (eGFR) less than 30 mL/min, QTc interval greater than 480 msec at baseline, uncorrected hypokalemia, coadministration with strong CYP3A4 inducers or strong CYP3A4 inhibitors
Required Medical Information	Diagnosis of covered use, submission of platelet count, serum potassium level, eGFR, and QTc interval, Child-Pugh score, documentation from a physical exam patient has splenomegaly.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VOQUEZNA (vonoprazan)

## Products Affected

- VOQUEZNA ORAL TABLET 10 MG, 20 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For erosive esophagitis and non-erosive gastroesophageal reflux disease (GERD), submission of current or previous therapies used to treat the condition (see Other Criteria). For Helicobacter pylori infection only, attestation patient will be administering with amoxicillin or a combination of amoxicillin and clarithromycin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	Up to 32 weeks based on covered use (see Other Criteria)
Other Criteria	For initial authorization for erosive esophagitis and non-erosive GERD, the patient must have tried and failed to have an adequate response to two different proton pump inhibitors or have contraindications to the proton pump inhibitor class. For non-erosive GERD, the initial coverage duration will be 4 weeks, with the option to reauthorize for an additional 20 weeks (for a total of 24 weeks of therapy per 365 days) if the provider attests to medical need. For H. pylori infection, a maximum of one 14-day course of vonoprazan will be approved. For all other indications, a maximum of one 32-week course of vonoprazan will be allowed per 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	Yes

# VORANIGO (vorasidenib)

---

## Products Affected

- VORANIGO ORAL TABLET 10 MG, 40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of IDH1 or IDH2 mutation, pregnancy status for female patients of childbearing potential, attestation patient has had at least one prior surgery (biopsy, sub-total resection, or gross total resection).
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	6 months
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# VOWST (fecal microbiota spores, live-brpk) EGWP

## Products Affected

- VOWST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use with the requirement patient is being treated after at least 2 recurrent (3 total) Clostridioides difficile infections (confirmation of pathogen with stool test or other confirmatory test), submission of time of last planned dose of antibiotic for latest recurrent C. difficile infection and attestation patient will be using a bowel cleanse the evening prior to starting Vowst.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 course (3 days)
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# WELIREG (belzutifan)

## Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For von Hippel-Lindau (VHL) disease, confirmation of a germline VHL alteration and attestation patient does not require immediate surgery. For advanced renal cell carcinoma, confirmation patient was previously treated with a programmed death receptor-1 or programmed death-ligand 1 inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

## XALKORI (crizotinib)

### Products Affected

- XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming tumor is ALK or ROS1-positive, pregnancy status for female patients of childbearing potential.
Age Restrictions	For ALK-positive systemic anaplastic large cell lymphoma only, 1 year of age to 21 years of age
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



# XDEMVY (lotilaner)

---

## Products Affected

- XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use including documentation of presence of mites upon examination of eyelashes by light microscopy or presence of collarettes on slit lamp examination, documentation of at least mild erythema of lid margin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to ophthalmology and optometry
Coverage Duration	6 weeks
Other Criteria	The safety and efficacy of retreating with additional courses has not been fully described. For this reason, only one 6-week treatment course will be allowed every 365 days.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# XOSPATA (gilteritinib)

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia or hypomagnesemia, coadministration with dual strong CYP3A/P-glycoprotein inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of FLT3 mutation, baseline serum potassium and magnesium levels, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# XURIDEN (uridine triacetate)

---

## Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline CBC including neutrophil count and mean corpuscular volume, baseline urine orotic acid level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For each annual reauthorization, documentation of improvements or stabilization of urine orotic acid level, neutrophil count, and mean corpuscular volume is required.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No

# ZYKADIA (ceritinib)

---

## Products Affected

- ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No
Prerequisite Therapy Required	No



## Index

ACTIMMUNE .....	2	<i>carglumic acid oral tablet soluble</i> .....	30
<i>adapalene external cream</i> .....	207	CERDELGA .....	31
<i>adapalene external gel 0.3 %</i> .....	207	CHENODAL .....	34
ADEMPAS ORAL TABLET 0.5 MG, 1 MG, 1.5 MG, 2 MG, 2.5 MG .....	3	CHOLBAM .....	35
AIMOVIG SUBCUTANEOUS SOLUTION AUTO-INJECTOR 140 MG/ML, 70 MG/ML .....	33	COBENFY .....	177
AJOVY .....	33	COBENFY STARTER PACK .....	177
AKEEGA .....	157	COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG .....	36
ALECENSA .....	4	COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG .....	36
ALUNBRIG .....	5	COMETRIQ (60 MG DAILY DOSE) .....	36
ALYQ .....	158	COPIKTRA ORAL CAPSULE 15 MG, 25 MG .....	37
<i>ambrisentan oral tablet 10 mg, 5 mg</i> .....	158	CORLANOR ORAL SOLUTION .....	154
<i>apomorphine hcl subcutaneous</i> .....	144	CORTROPHIN .....	38
AQNEURSA .....	6	CORTROPHIN GEL .....	38
ARIKAYCE .....	7	COTELLIC .....	39
ASCOMP-CODEINE .....	22	CRESEMBA ORAL CAPSULE 186 MG .....	40
AUGTYRO .....	8	CYSTADROPS .....	41
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG .....	9	CYSTARAN .....	41
AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HOUR 12 MG, 18 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG .....	9	<i>dalfampridine er</i> .....	42
AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG .....	9	DANZITEN .....	125
AUVELITY .....	10	<i>dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg</i> .....	43
AVMAPKI FAKZYNJA CO-PACK .....	11	DAURISMO ORAL TABLET 100 MG, 25 MG .....	44
AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG .....	12	<i>deferasirox oral tablet soluble</i> .....	45
BALVERSA .....	13	<i>deferiprone</i> .....	46
BENLYSTA SUBCUTANEOUS .....	15	DIACOMIT .....	47
BESREMI .....	16	<i>dichlorphenamide</i> .....	48
<i>bexarotene external</i> .....	17	<i>diclofenac epolamine external</i> .....	49
<i>bosentan oral tablet 125 mg, 62.5 mg</i> .....	158	<i>diclofenac sodium external gel 3 %</i> .....	154
BOSULIF ORAL CAPSULE 100 MG, 50 MG .....	19	<i>digoxin oral tablet 250 mcg</i> .....	50
BOSULIF ORAL TABLET .....	19	DOPTLET ORAL TABLET 20 MG .....	51
BRAFTOVI ORAL CAPSULE 75 MG .....	20	<i>dronabinol</i> .....	52
BRUKINSA .....	21	DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR .....	53
<i>butalbital-acetaminophen oral tablet 50-300 mg, 50-325 mg</i> .....	22	DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 200 MG/1.14ML, 300 MG/2ML .....	53
<i>butalbital-apap-caff-cod</i> .....	22	<i>eltrombopag olamine oral packet</i> .....	55
<i>butalbital-apap-caffeine oral capsule</i> .....	22	<i>eltrombopag olamine oral tablet 12.5 mg, 25 mg, 50 mg, 75 mg</i> .....	55
<i>butalbital-apap-caffeine oral tablet 50-325-40 mg</i> ....	22	EMGALITY .....	33
<i>butalbital-asa-caff-codeine</i> .....	22	EMGALITY (300 MG DOSE) .....	33
<i>butalbital-aspirin-caffeine oral capsule</i> .....	22	ENSACOVE .....	56
BYLVAY .....	24	EOHILIA .....	57
CABLIVI .....	25	EPIDIOLEX .....	58
CABOMETYX .....	26	ERIVEDGE .....	59
CALQUENCE ORAL TABLET .....	27	ERLEADA .....	157
CAMZYOS .....	28	<i>eszopiclone</i> .....	178
CAPLYTA ORAL CAPSULE 10.5 MG, 21 MG, 42 MG ...	177	<i>everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg</i> ..	60
CAPRELSA .....	29	<i>everolimus oral tablet soluble</i> .....	60
		EVRYSDI .....	61
		<i>ezetimibe-simvastatin oral tablet 10-80 mg</i> .....	180

FABHALTA .....	62	KINERET SUBCUTANEOUS SOLUTION PREFILLED	
FANAPT ORAL TABLET 1 MG, 10 MG, 12 MG, 2 MG, 4 MG, 6 MG, 8 MG .....	177	SYRINGE .....	18
FANAPT TITRATION PACK .....	177	KISQALI (200 MG DOSE) .....	97
FANAPT TITRATION PACK A .....	177	KISQALI (400 MG DOSE) .....	97
FANAPT TITRATION PACK B ORAL TABLET .....	177	KISQALI (600 MG DOSE) .....	97
FANAPT TITRATION PACK C ORAL TABLET .....	177	KOSELUGO .....	98
FILSPARI .....	63	KRAZATI .....	99
FINTEPLA .....	64	<i>lapatinib ditosylate</i> .....	100
FIRDAPSE .....	65	LAZCLUZE ORAL TABLET 240 MG, 80 MG .....	101
FOTIVDA .....	66	<i>lenalidomide</i> .....	102
FRUZAQLA ORAL CAPSULE 1 MG, 5 MG .....	67	LENVIMA (10 MG DAILY DOSE) .....	103
GAMMAGARD .....	84	LENVIMA (12 MG DAILY DOSE) .....	103
GAMMAGARD S/D LESS IGA .....	84	LENVIMA (14 MG DAILY DOSE) .....	103
GATTEX .....	68	LENVIMA (18 MG DAILY DOSE) .....	103
GAVRETO .....	69	LENVIMA (20 MG DAILY DOSE) .....	103
<i>gefitinib</i> .....	70	LENVIMA (24 MG DAILY DOSE) .....	103
GILOTRIF .....	71	LENVIMA (4 MG DAILY DOSE) .....	103
GOMEKLI .....	73	LENVIMA (8 MG DAILY DOSE) .....	103
HAEGARDA .....	76	LEUKINE INJECTION SOLUTION RECONSTITUTED .....	104
HERNEXEOS .....	77	<i>l-glutamine oral packet</i> .....	105
IBRANCE .....	78	<i>lidocaine external patch 5 %</i> .....	106
IBTROZI .....	79	LIDOCAN .....	106
<i>icatibant acetate subcutaneous solution prefilled</i> <i>syringe</i> .....	75	LIVTENCITY .....	107
ICLUSIG .....	80	LODOCO .....	108
IDHIFA .....	81	LONSURF .....	109
IMBRUVICA ORAL CAPSULE .....	82	LORBRENA ORAL TABLET 100 MG, 25 MG .....	110
IMBRUVICA ORAL SUSPENSION .....	82	LUMAKRAS .....	111
IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG .....	82	LYBALVI .....	112
<i>imkeldi</i> .....	83	LYNPARZA ORAL TABLET .....	113
INBRIJA .....	144	LYTGOBI (12 MG DAILY DOSE) .....	114
INCRELEX .....	116	LYTGOBI (16 MG DAILY DOSE) .....	114
INLYTA .....	85	LYTGOBI (20 MG DAILY DOSE) .....	114
INQOVI .....	86	MAVYRET .....	115
INREBIC .....	87	<i>megestrol acetate oral suspension 40 mg/ml, 625</i> <i>mg/5ml</i> .....	117
INVEGA HAFYERA .....	88	MEKINIST .....	118
INVEGA TRINZA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 273 MG/0.88ML, 410 MG/1.32ML, 546 MG/1.75ML, 819 MG/2.63ML .....	88	MEKTOVI .....	20
ITOVEBI ORAL TABLET 3 MG, 9 MG .....	89	<i>memantine hcl-donepezil hcl</i> .....	119
<i>ivabradine hcl oral tablet 5 mg, 7.5 mg</i> .....	154	<i>mifepristone oral tablet 300 mg</i> .....	120
IWILFIN .....	90	<i>miglustat</i> .....	121
JAKAFI .....	91	MODEYSO .....	122
JAVYGTOR .....	175	MOUNJARO SUBCUTANEOUS SOLUTION AUTO- INJECTOR 10 MG/0.5ML, 12.5 MG/0.5ML, 15 MG/0.5ML, 2.5 MG/0.5ML, 5 MG/0.5ML, 7.5 MG/0.5ML .....	72
JAYPIRCA .....	92	MYTESI .....	123
JOENJA .....	93	NAMZARIC ORAL CAPSULE EXTENDED RELEASE 24 HOUR 7-10 MG .....	119
JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG .....	94	NAYZILAM .....	154
KALYDECO .....	32	NERLYNX .....	124
KERENDIA .....	95	NEXLETOL .....	14
<i>ketoconazole oral</i> .....	96	NEXLIZET .....	14
		<i>nilotinib hcl</i> .....	125

NINLARO .....	126	PIQRAY (200 MG DAILY DOSE) .....	149
<i>nitisinone</i> .....	127	PIQRAY (250 MG DAILY DOSE) .....	149
NORDITROPIN FLEXPOR SUBCUTANEOUS SOLUTION		PIQRAY (300 MG DAILY DOSE) .....	149
PEN-INJECTOR .....	74	<i>pirfenidone oral tablet 267 mg, 801 mg</i> .....	150
NUBEQA .....	157	POMALYST .....	151
NUCALA SUBCUTANEOUS SOLUTION AUTO-		<i>pretomanid</i> .....	152
INJECTOR .....	128	PREVYMIS ORAL TABLET .....	153
NUCALA SUBCUTANEOUS SOLUTION PREFILLED		PRIVIGEN .....	84
SYRINGE 100 MG/ML .....	128	PROLASTIN-C INTRAVENOUS SOLUTION .....	155
NUCALA SUBCUTANEOUS SOLUTION		<i>promethazine hcl oral solution 6.25 mg/5ml</i> .....	156
RECONSTITUTED .....	128	<i>promethazine hcl oral tablet</i> .....	156
NUEDEXTA .....	130	<i>promethazine hcl rectal suppository 12.5 mg, 25 mg</i> .....	156
NUPLAZID ORAL CAPSULE .....	131	<i>promethazine-phenylephrine</i> .....	156
NUPLAZID ORAL TABLET 10 MG .....	131	PROMETHEGAN RECTAL SUPPOSITORY 25 MG, 50	
NURTEC .....	33	MG .....	156
OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML,		QINLOCK .....	159
10 GM/100ML, 10 GM/200ML, 2 GM/20ML, 2.5		QULIPTA .....	33
GM/50ML, 20 GM/200ML, 5 GM/100ML, 5		RADICAVA ORS .....	160
GM/50ML .....	84	RADICAVA ORS STARTER KIT .....	160
ODOMZO .....	132	RECORLEV .....	161
OFEV .....	133	REPATHA .....	162
OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG ...	134	REPATHA PUSHTRONEX SYSTEM .....	162
OJEMDA ORAL SUSPENSION RECONSTITUTED .....	135	REPATHA SURECLICK .....	162
OJEMDA ORAL TABLET 100 MG, 100 MG (16 PACK),		RETACRIT INJECTION SOLUTION 10000 UNIT/ML,	
100 MG (24 PACK) .....	135	2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML,	
OJJAARA ORAL TABLET 100 MG, 150 MG, 200 MG ...	136	4000 UNIT/ML, 40000 UNIT/ML .....	163
ONUREG .....	137	RETEVMO ORAL TABLET 120 MG, 160 MG, 40 MG,	
OPSUMIT .....	158	80 MG .....	164
ORENITRAM .....	158	REVCovi .....	165
ORENITRAM MONTH 1 .....	158	REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG ..	166
ORENITRAM MONTH 2 .....	158	REXULTI ORAL TABLET 0.25 MG, 0.5 MG, 1 MG, 2	
ORENITRAM MONTH 3 .....	158	MG, 3 MG, 4 MG .....	177
ORIAHNN .....	138	REZDIFFRA .....	167
ORILISSA ORAL TABLET 150 MG, 200 MG .....	139	REZLIDHIA .....	168
ORKAMBI .....	32	REZUROCK .....	169
ORMALVI .....	48	RIVFLOZA .....	170
ORSERDU .....	140	ROMVIMZA .....	171
OTEZLA ORAL TABLET 30 MG .....	18	ROZLYTREK .....	172
OTEZLA ORAL TABLET THERAPY PACK 10 & 20 & 30		RUBRACA .....	173
MG .....	18	RYBELSUS (FORMULATION R2) ORAL TABLET 1.5	
OXERVATE .....	141	MG, 4 MG, 9 MG .....	72
OZEMPIC (0.25 OR 0.5 MG/DOSE) SUBCUTANEOUS		RYBELSUS ORAL TABLET 14 MG, 3 MG, 7 MG .....	72
SOLUTION PEN-INJECTOR 2 MG/3ML .....	72	RYDAPT .....	174
OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION		SAJAZIR SUBCUTANEOUS SOLUTION PREFILLED	
PEN-INJECTOR 4 MG/3ML .....	72	SYRINGE .....	75
OZEMPIC (2 MG/DOSE) .....	72	<i>sapropterin dihydrochloride oral packet</i> .....	175
PANRETIN .....	143	<i>sapropterin dihydrochloride oral tablet</i> .....	175
<i>pazopanib hcl</i> .....	145	SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG ....	176
PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML	154	SECUADO TRANSDERMAL PATCH 24 HOUR 3.8	
PEGASYS SUBCUTANEOUS SOLUTION PREFILLED		MG/24HR, 5.7 MG/24HR, 7.6 MG/24HR .....	177
SYRINGE .....	154	SIGNIFOR .....	179
PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG ...	147	<i>sildenafil citrate oral suspension reconstituted</i> .....	158
PHEBURANE .....	148	<i>sildenafil citrate oral tablet 20 mg</i> .....	158



<i>simvastatin oral tablet 80 mg</i> .....	180	TRIKAFTA ORAL THERAPY PACK .....	32
SIRTURO .....	181	TRULICITY SUBCUTANEOUS SOLUTION AUTO- INJECTOR .....	72
SIVEXTRO .....	182	TRUQAP ORAL TABLET .....	208
SKYCLARYS .....	183	TRUQAP ORAL TABLET THERAPY PACK .....	208
<i>sodium oxybate</i> .....	142	TUKYSA ORAL TABLET 150 MG, 50 MG .....	209
<i>sofosbuvir-velpatasvir</i> .....	184	TURALIO ORAL CAPSULE 125 MG .....	210
SOHONOS .....	185	TYENNE SUBCUTANEOUS .....	18
SOMAVERT .....	186	UBRELVY .....	33
<i>sorafenib tosylate</i> .....	187	UDENYCA .....	146
SOTYKTU .....	18	UDENYCA ONBODY .....	146
STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML ..	18	<i>ustekinumab subcutaneous</i> .....	18
STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	18	UZEDY .....	211
STIVARGA .....	188	VALCHLOR .....	212
SUCRAID .....	189	VALTOCO 10 MG DOSE .....	154
<i>sunitinib malate</i> .....	190	VALTOCO 15 MG DOSE NASAL LIQUID THERAPY PACK 2 X 7.5 MG/0.1ML .....	154
SUNOSI ORAL TABLET 150 MG, 75 MG .....	191	VALTOCO 20 MG DOSE NASAL LIQUID THERAPY PACK 2 X 10 MG/0.1ML .....	154
SYMDEKO .....	32	VALTOCO 5 MG DOSE .....	154
TABRECTA .....	192	VANFLYTA .....	213
<i>tadalafil (pah)</i> .....	158	VANRAFIA .....	214
<i>tadalafil oral tablet 2.5 mg, 5 mg</i> .....	193	VELSIPITY .....	18
TAFINLAR .....	118	VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG ..	215
TAGRISSO .....	195	VENCLEXTA STARTING PACK .....	215
TAKHZYRO SUBCUTANEOUS SOLUTION .....	76	VEOZAH .....	216
TAKHZYRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 300 MG/2ML .....	76	VERQUVO ORAL TABLET 10 MG, 2.5 MG, 5 MG .....	217
TALZENNA ORAL CAPSULE 0.1 MG, 0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG .....	196	VERZENIO .....	218
<i>tasimelteon</i> .....	197	VIBERZI .....	219
TAVNEOS .....	198	VIJOICE ORAL PACKET .....	220
<i>tazarotene external cream</i> .....	207	VIJOICE ORAL TABLET THERAPY PACK 125 MG, 200 & 50 MG, 50 MG .....	220
<i>tazarotene external gel</i> .....	207	VITRAKVI ORAL CAPSULE 100 MG, 25 MG .....	221
TAZVERIK .....	199	VITRAKVI ORAL SOLUTION .....	221
TENCON ORAL TABLET 50-325 MG .....	22	VIVJOA .....	222
TEPMETKO .....	200	VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG .....	223
<i>teriparatide subcutaneous solution pen-injector 560 mcg/2.24ml, 620 mcg/2.48ml</i> .....	201	VONJO .....	224
<i>testosterone transdermal gel 10 mg/act (2%), 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)</i> .....	203	VOQUEZNA ORAL TABLET 10 MG, 20 MG .....	225
<i>testosterone transdermal solution</i> .....	203	VORANIGO ORAL TABLET 10 MG, 40 MG .....	226
TIBSOVO .....	204	VOWST .....	227
<i>tolvaptan oral tablet 15 mg, 30 mg</i> .....	205	VOYDEYA ORAL TABLET .....	228
<i>tolvaptan oral tablet 15 mgtolvaptan (hyponatremia), 30 mgtolvaptan (hyponatremia) ...</i>	206	VOYDEYA ORAL TABLET THERAPY PACK .....	228
<i>tolvaptan oral tablet therapy pack 15 mg, 30 &amp; 15 mg, 45 &amp; 15 mg, 60 &amp; 30 mg, 90 &amp; 30 mg</i> .....	205	VYNDAMAX .....	194
<i>tretinoin external cream</i> .....	207	VYNDAQEL .....	194
<i>tretinoin external gel 0.01 %, 0.025 %</i> .....	207	WEGOVY SUBCUTANEOUS SOLUTION AUTO- INJECTOR 0.25 MG/0.5ML, 0.5 MG/0.5ML, 1 MG/0.5ML, 1.7 MG/0.75ML, 2.4 MG/0.75ML .....	229
TRIDACAINE II .....	106	WELIREG .....	231
TRIKAFTA ORAL TABLET THERAPY PACK 100-50-75 & 150 MG, 50-25-37.5 & 75 MG .....	32	WINREVAIR SUBCUTANEOUS KIT 2 X 45 MG, 2 X 60 MG, 45 MG, 60 MG .....	232
		XALKORI .....	233
		XDEMVI .....	234
		XERMELO .....	235

XIFAXAN ORAL TABLET 550 MG .....	236
XOLAIR .....	237
XOSPATA .....	239
XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG .....	240
XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG .....	240
XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG .....	240
XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG .....	240
XPOVIO (60 MG TWICE WEEKLY) .....	240
XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG .....	240
XPOVIO (80 MG TWICE WEEKLY) .....	240
XTANDI .....	157
XURIDEN .....	241
YARGESA .....	121
<i>zaleplon</i> .....	178
ZEJULA ORAL TABLET .....	242
ZELBORAF .....	39
ZILBRYSQ .....	243
<i>zolpidem tartrate oral tablet</i> .....	178
ZORYVE EXTERNAL CREAM 0.3 % .....	244
ZTALMY .....	245
ZURZUVAE .....	246
ZYDELIG .....	247
ZYKADIA ORAL TABLET .....	248