

## PRIOR AUTHORIZATION CRITERIA

This list is current as of October 1, 2021, and pertains to the following formularies:

2022 Pharmacy Benefit Dimensions Prescription Drug Plan (PDP) Part D 3 Tier Formulary	Version 8
--	-----------

Pharmacy Benefit Dimensions requires you (or your physician) to get prior authorization for certain drugs listed on the formularies above. This means that you will need to get approval from us before you fill your prescriptions. If you do not get approval, we may not cover the drug. These drugs are listed with a “PA” in the Requirements/Notes column on the formularies. This document contains the Prior Authorization requirements that are associated with the formularies listed above.

If you have any questions, please contact our Medicare Member Services Department at 1-800-667-5936 or, for TTY 711, October 1<sup>st</sup> – March 31<sup>st</sup>: Monday through Sunday from 8 a.m. to 8 p.m. ET, April 1<sup>st</sup> – September 30<sup>th</sup>: Monday through Friday from 8 a.m. to 8 p.m. ET.

Pharmacy Benefit Dimensions is a subsidiary of Independent Health. Independent Health is a PDP with a Medicare contract. Enrollment in Pharmacy Benefit Dimensions PDP depends on contract renewal between Independent Health and CMS.

The formulary may change at any time. You will receive notice when necessary.

# ABILIFY MYCITE (aripiprazole with sensor)

---

## Products Affected

- ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET 10 MG, 15 MG, 2 MG, 20 MG, 30 MG, 5 MG
- ABILIFY MYCITE STARTER KIT ORAL TABLET 10 MG, 15 MG, 2 MG, 20 MG, 30 MG, 5 MG
- ABILIFY MYCITE ORAL TABLET 10 MG, 15 MG, 2 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of previous aripiprazole use (see Other Criteria).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval, the patient must have documentation of at least a one-month trial of generic aripiprazole solution, tablets, or orally-disintegrating tablets.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ACTHAR (corticotropin)

## Products Affected

- ACTHAR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Request for IV administration, treatment of patients under 2 years of age in whom congenital infections are suspected, patients with scleroderma, osteoporosis, systemic fungal infections, ocular herpes simplex, recent surgery, a history of or presence of a peptic ulcer, congestive heart failure, uncontrolled hypertension, primary adrenocortical insufficiency, adrenocortical hyperfunction, or sensitivity to proteins of porcine origin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of blood pressure reading and baseline serum sodium and potassium levels.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated blood pressure, sodium, and potassium levels since the previous authorization will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ACTIMMUNE (interferon gamma-1b)

---

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ADEMPAS (riociguat)

---

## Products Affected

- ADEMPAS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe (Child-Pugh class C) hepatic impairment, creatinine clearance below 15 mL/min or on dialysis, concurrent use with nitrates or nitric oxide donors in any form, concurrent use with phosphodiesterase inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of negative pregnancy test result for female patients of childbearing age, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# AKYNZEO (netupitant/palonosetron)

---

## Products Affected

- AKYNZEO ORAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, severe renal impairment, end-stage renal disease
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation patient will receive concurrent dexamethasone therapy as indicated based on level of chemotherapy regimen emetogenicity.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. If the medication is being administered related to cancer treatment and is a full replacement for intravenous administration of antiemetic therapy within 48 hours of cancer treatment, it is covered as a Part B benefit. To be eligible for Part B coverage, the prescribing physician must indicate this on the prescription. Otherwise it may be covered as a Part D benefit.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ALECENSA (alectinib)

---

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of ALK-positive tumor.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ALPHA-1-PROTEINASE INHIBITORS

## Products Affected

- ARALAST NP INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG, 500 MG
- GLASSIA
- PROLASTIN-C INTRAVENOUS SOLUTION
- RECONSTITUTED ZEMAIRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Individuals with immunoglobulin A (IgA) deficiency who have known antibodies against IgA
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation that patient has clinically evident emphysema secondary to congenital alpha-1-PI deficiency by submission of pulmonary function testing (e.g., spirometry or body plethysmography), X-ray radiography, or diffusing capacity of the lung for carbon monoxide (DLCO).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. Continuation of therapy requests require objective documentation from the prescriber that the patient's symptoms have improved. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# ALUNBRIG (brigatinib)

---

## Products Affected

- ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of ALK-positive tumor.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# AMANTADINE EXTENDED-RELEASE PRODUCTS

---

## Products Affected

- GOCOVRI

PA Criteria	Criteria Details
Exclusion Criteria	End stage renal disease (creatinine clearance below 15 mL/min)
Required Medical Information	Diagnosis of covered use, documentation patient tried and failed immediate-release amantadine.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ANADROL-50 (oxymetholone)

---

## Products Affected

- ANADROL-50

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Carcinoma of the prostate or breast in male patients, carcinoma of the breast in females with hypercalcemia, women who are or may become pregnant, nephrosis or the nephrotic phase of nephritis, severe hepatic dysfunction
<b>Required Medical Information</b>	Diagnosis of covered use, submission of CBC, liver function tests, and pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	PA applies to all. Updated CBC since the previous authorization will be required for reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ARANESP (darbepoetin alfa)

## Products Affected

- ARANESP (ALBUMIN FREE) INJECTION SOLUTION 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 300 MCG/ML, 40 MCG/ML, 60 MCG/ML
- ARANESP (ALBUMIN FREE) INJECTION SOLUTION

PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of hemoglobin or hematocrit level, serum iron, total iron-binding capacity (TIBC), and transferrin within 30 days of request date, documentation that the patient does not have uncontrolled hypertension.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For non-ESRD-related conditions: 90 days. For ESRD-related conditions: 1 year.
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ARCALYST (rilonacept)

---

## Products Affected

- ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	Active or chronic infection, coadministration with TNF-blocking agents
Required Medical Information	Diagnosis of covered use, TB skin test result obtained within the past 12 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ARIKAYCE (amikacin inhalation)

---

## Products Affected

- ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	Non-refractory Mycobacterium avium complex (MAC) lung disease
Required Medical Information	Diagnosis of covered use, submission of other therapies that have been tried and failed or cannot be used because of a contraindication.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	This medication is covered as a Part B benefit except for enrollees residing in a long-term care facility. PA applies to all when covered as a Part D benefit.
Indications	All Medically-accepted Indications.
Off Label Uses	

# AURYXIA (ferric citrate)

---

## Products Affected

- AURYXIA

PA Criteria	Criteria Details
Exclusion Criteria	Iron overload syndrome
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# BALVERSA (erdafitinib)

---

## Products Affected

- BALVERSA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP2C9 or CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of FDA-approved companion test showing susceptible FGFR2 or FGFR3 genetic alterations, prior chemotherapy regimen(s) used.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# BEMPEDOIC ACID

---

## Products Affected

- NEXLETOL
- NEXLIZET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant pravastatin utilization with doses above 40 mg/day, concomitant simvastatin utilization with doses above 20 mg/day
<b>Required Medical Information</b>	Diagnosis of covered use, submission of current or previous lipid-lowering therapies.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval, the patient must currently be using a statin plus ezetimibe or the patient must have tried and failed to have an adequate response to or had an intolerance to at least two statins or one statin and ezetimibe. At least one statin previously tried and failed must be a hydrophilic statin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# BENLYSTA (belimumab)

---

**Products Affected**

- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe active central nervous system lupus, patients using other biologic medications or intravenous cyclophosphamide
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation that the patient is taking standard therapy defined as at least one of the following: corticosteroids, NSAIDs, antimalarials, or immunosuppressants.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# BIOLOGIC RESPONSE MODIFIERS

## Products Affected

- ACTEMRA ACTPEN
- ACTEMRA SUBCUTANEOUS
- CIMZIA PREFILLED
- CIMZIA STARTER KIT
- CIMZIA SUBCUTANEOUS KIT 2 X 200 MG
- KEVZARA
- OTEZLA
- SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- TREMFYA
- ZEPOSIA
- ZEPOSIA 7-DAY STARTER PACK
- ZEPOSIA STARTER KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	For Zeposia for the treatment of multiple sclerosis, only diagnosis of covered use is required. For all other drugs managed by this policy and for Zeposia for indications other than multiple sclerosis, diagnosis of covered use, submission of previous therapies. For all drugs managed by this policy except Otezla and Zeposia, submission of baseline latent tuberculosis screening test (Mantoux tuberculin skin test [a.k.a. PPD test] or interferon-gamma release assay [IGRA]).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. With the exception of Zeposia for the treatment of multiple sclerosis only, for approval of a drug managed by this policy, the patient must have tried and failed to have an adequate response to or had an intolerance to at least two preferred agents (Cosentyx, Enbrel, Humira, Rinvoq, Skyrizi, Stelara, and Xeljanz/Xeljanz XR) for the indication submitted, where possible. For all drugs managed by this policy except Otezla and Zeposia, if TB screening test returns a positive result, coverage will be delayed until latent TB is treated. For re-authorization, yearly TB screening test or chest X-ray required for patients who live in, work in, or travel to areas where TB exposure is likely while on treatment or for those who have previously had a positive TB screening test.
Indications	All Medically-accepted Indications.
Off Label Uses	

# BOSULIF (bosutinib)

---

## Products Affected

- BOSULIF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inhibitors or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of renal function testing. For accelerated or blast phase Ph+ CML, documentation of resistance or intolerance to at least one of the following prior therapies: imatinib, dasatinib, or nilotinib.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# BRAFTOVI/MEKTOVI (encorafenib/binimetinib)

---

## Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG
- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of BRAF V600E or V600K mutation, serum potassium, and serum magnesium. For metastatic melanoma, confirmation that encorafenib and binimetinib will be co-administered. For metastatic colorectal cancer, confirmation that encorafenib and cetuximab will be co-administered.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# BRONCHITOL (mannitol powder for inhalation)

---

## Products Affected

- BRONCHITOL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Documented Bronchitol Tolerance Test failure
<b>Required Medical Information</b>	Diagnosis of covered use, documentation patient has passed the Bronchitol Tolerance Test, attestation patient will not be using in combination with hypertonic (7%) sodium chloride nebulized solution.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval, the patient must must have tried and failed to have an adequate response to or had an intolerance to hypertonic (7%) sodium chloride nebulized solution.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# BRUKINSA (zanubrutinib)

---

## Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of prior chemotherapy regimen(s) used.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# BUTALBITAL-CONTAINING PRODUCTS IN OLDER PATIENTS

## Products Affected

- *ascomp-codeine*
- BUPAP ORAL TABLET 50-300 MG
- BUTALBITAL-ACETAMINOPHEN ORAL TABLET 50-300 MG, 50-325 MG
- BUTALBITAL-APAP-CAFF-COD
- *butalbital-apap-caffeine oral capsule*
- *butalbital-apap-caffeine oral tablet 50-325-40 mg*
- *butalbital-asa-caff-codeine*
- *butalbital-aspirin-caffeine oral capsule*
- TENCON ORAL TABLET 50-325 MG
- VTOL LQ
- ZEBUTAL ORAL CAPSULE 50-325-40 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation patient has tried and failed a preferred alternative such as ibuprofen or rizatriptan, or has contraindications to all alternatives.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	



# BYLVAY (odevixibat)

---

## Products Affected

- BYLVAY
- BYLVAY (PELLETS)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of liver transplant, clinical evidence of decompensated cirrhosis
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by molecular genetic testing, attestation drug-induced pruritus has been ruled out.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to gastroenterology and hepatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Attestation of improvement in pruritus symptoms and submission of liver function testing, including serum bilirubin, since the previous authorization will be required for subsequent reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# CABLIVI (caplacizumab-yhdp)

---

## Products Affected

- CABLIVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation drug will be given with plasma exchange and immunosuppressive therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology, hematology, and immunology
Coverage Duration	3 months
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# CABOMETYX (cabozantinib)

---

## Products Affected

- CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# CALQUENCE (acalabrutinib)

---

## Products Affected

- CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment, coadministration with strong CYP3A inhibitors or proton pump inhibitors
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# CAPLYTA (lumateperone)

---

## Products Affected

- CAPLYTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis, moderate or severe hepatic impairment, coadministration with moderate or strong CYP3A4 inhibitors or CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of previous therapies used for indication and liver function testing or Child-Pugh score.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval, the patient must have tried and failed to have an adequate response to or had an intolerance to aripiprazole and at least one other generic second-generation atypical antipsychotic (e.g., paliperidone, quetiapine, risperidone, etc.) or Latuda.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# CAPRELSA (vandetanib)

---

**Products Affected**

- CAPRELSA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of congenital long QT syndrome, Torsades de pointes, uncompensated heart failure, or bradyarrhythmias, moderate or severe hepatic impairment, QTcF interval greater than 450 msec
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline serum potassium, calcium, magnesium, ALT, AST, bilirubin, TSH, creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), ECG, and pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# CARBAGLU (carglumic acid)

---

## Products Affected

- CARBAGLU

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of plasma ammonia level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated plasma ammonia level since the previous authorization will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# CERDELGA (eliglustat)

---

## Products Affected

- CERDELGA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pre-existing cardiac disease, long QT syndrome, coadministration with Class Ia or Class III antiarrhythmics, patients who are extensive or intermediate CYP2D6 metabolizers taking a strong CYP2D6 inhibitor with a strong or moderate CYP3A inhibitor, intermediate and poor CYP2D6 metabolizers taking a strong CYP3A inhibitor
<b>Required Medical Information</b>	Diagnosis of covered use, submission of CYP2D6 metabolizer status as detected by an FDA-cleared test for determining CYP2D6 genotype, liver function testing or Child-Pugh score.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated liver function testing or Child-Pugh score since the previous authorization will be required for subsequent reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# CHENODAL (chenodiol)

---

## Products Affected

- CHENODAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known hepatocyte dysfunction, bile duct abnormalities such as intrahepatic cholestasis, primary biliary cirrhosis, or sclerosing cholangitis, radiopaque stones, nonvisualizing gallbladder confirmed as nonvisualizing after 2 consecutive single doses of dye, compelling reasons for gallbladder surgery
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 months
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# CHOLBAM (cholic acid)

---

## Products Affected

- CHOLBAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of liver function testing.
Age Restrictions	
Prescriber Restrictions	Restricted to hepatology, gastroenterology, and pediatric gastroenterology
Coverage Duration	Initially 3 months, then 1 year
Other Criteria	PA applies to all. Documentation of liver function improvement without complete biliary obstruction or persistent clinical or laboratory indications of worsening liver function or cholestasis will be required for initial reauthorization after the first 3 months. Updated liver function testing since the previous authorization will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# COMETRIQ (cabozantinib)

---

## Products Affected

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	Severe hepatic impairment (Child-Pugh class C)
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# COPIKTRA (duvelisib)

---

## Products Affected

- COPIKTRA ORAL CAPSULE 15 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of at least two prior therapies tried and failed, submission of pregnancy status for female patients of childbearing potential, attestation patient will receive prophylaxis for <i>Pneumocystis jirovecii</i> pneumonia (PJP) and, if necessary, cytomegalovirus.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology or oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# CORLANOR (ivabradine)

## Products Affected

- CORLANOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute decompensated heart failure, clinically significant hypotension, clinically significant bradycardia, severe hepatic impairment, pacemaker dependence (heart rate maintained exclusively by the pacemaker), or sick sinus syndrome, sinoatrial block, or 3rd degree AV block unless a functioning demand pacemaker is present, coadministration with strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use described as is indicated (1) to reduce the risk of hospitalization for worsening heart failure in patients with stable, symptomatic chronic heart failure with left ventricular ejection fraction less than or equal to 35%, who are in sinus rhythm with resting heart rate greater than or equal to 70 beats per minute and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use or (2) for stable symptomatic heart failure due to dilated cardiomyopathy in patients who are in sinus rhythm with an elevated heart rate. Submission of current baseline blood pressure reading, confirmation that patient does not have any of the following: (1) acute decompensated heart failure, (2) sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present, (3) resting heart rate less than 60 bpm prior to treatment, (4) severe hepatic impairment, (5) pacemaker dependence (heart rate maintained exclusively by the pacemaker). For patients under 18 years old, (1) left ventricular ejection fraction less than or equal to 45% and (2) resting heart rate greater than or equal to the following age-stratified requirements: (a) 105 beats per minute in ages 6 to 12 months old, (b) 95 beats per minute in ages 1 to 3 years old, (c) 75 beats per minute in ages 3 to 5 years old, and (d) 70 beats per minute in ages 5 to 18 years old.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to cardiology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Continuation requires documentation of clinically relevant response to therapy.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# CYSTEAMINE EYE DROPS

---

## Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DALFAMPRIDINE

---

## Products Affected

- *dalfampridine er*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of seizure, moderate or severe renal impairment (CrCl less than or equal to 50 mL/min)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance) and objective measurement of walking speed, confirmation that patient is able to walk.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	Initially 3 months, then 1 year
<b>Other Criteria</b>	PA applies to all. Documentation the patient has demonstrated an improvement in walking speed from baseline measure (or maintenance of improvement if patient has been on long-term therapy) since starting medication will be required for initial reauthorization after the first 3 months. Updated creatinine clearance since the previous authorization and confirmation patient is able to walk will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# DAURISMO (glasdegib)

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, confirmation patient will also be receiving cytarabine as part of chemotherapeutic regimen. If patient is under 75 years of age, documentation of comorbidities that preclude use of intensive induction chemotherapy, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# DAYVIGO (lemborexant)

---

## Products Affected

- DAYVIGO ORAL TABLET 10 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	Narcolepsy
Required Medical Information	Diagnosis of covered use. Patient must have tried and failed to tolerate or had an inadequate response to two covered alternative therapies recommended by the American Academy of Sleep Medicine (doxepin, eszopiclone, ramelteon, suvorexant, temazepam, zaleplon, zolpidem) including one non-suvorexant therapy for sleep maintenance (doxepin, eszopiclone, temazepam) if that is the diagnosis of covered use.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DEFERASIROX

## Products Affected

- *deferasirox granules*
- *deferasirox oral tablet*
- *deferasirox oral tablet soluble*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, estimated glomerular filtration rate less than 40 mL per min, platelet count below $50 \times 10^9/L$ , high-risk myelodysplastic syndromes, advanced malignancies
<b>Required Medical Information</b>	Diagnosis of covered use, submission of CBC, LFTs, serum creatinine, ferritin, and urine protein values, estimated glomerular filtration rate, documentation that member has had yearly ophthalmic and auditory testing.
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	PA applies to all. Updated ferritin level within last 3 months and updated CBC, LFT, urine protein value, estimated glomerular filtration rate, and ophthalmic and auditory testing since the previous authorization will be required for subsequent reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# DEFERIPRONE

---

## Products Affected

- *deferiprone*
- FERRIPROX ORAL SOLUTION
- FERRIPROX ORAL TABLET 1000 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of serum ferritin levels, CBC, ANC, platelet count, and serum ALT.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DIACOMIT (stiripentol)

---

## Products Affected

- DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe renal impairment, moderate or severe hepatic impairment
Required Medical Information	Diagnosis of covered use, confirmation patient is also receiving clobazam.
Age Restrictions	2 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only. Monotherapy requests for Dravet syndrome will not be approved as there are no clinical data to support using stiripentol in this manner.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DICLOFENAC 1% GEL

---

## Products Affected

- *diclofenac sodium external gel*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Use during the peri-operative period in the setting of coronary artery bypass graft (CABG) surgery
<b>Required Medical Information</b>	Diagnosis of covered use defined as the relief of pain of osteoarthritis of joints amenable to topical treatment, such as the knees and hands.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Not evaluated for use on joints of the spine, hip, or shoulder and therefore requests for use on these areas will not be approved.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

# DICLOFENAC 1.5% TOPICAL SOLUTION

---

## Products Affected

- *diclofenac sodium external solution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Use during the peri-operative period in the setting of coronary artery bypass graft (CABG) surgery
<b>Required Medical Information</b>	Diagnosis of covered use defined as the relief of pain of osteoarthritis of the knees.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. FDA-approved only for use on knee joints and therefore requests for other uses will not be approved.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

# DICLOFENAC 3% GEL

---

## Products Affected

- *diclofenac sodium external gel*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology
Coverage Duration	90 days
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DICLOFENAC PATCH

---

## Products Affected

- *diclofenac epolamine external*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery, use on non-intact or damaged skin resulting from any etiology including exudative dermatitis, eczema, infection lesions, burns, or wounds, pregnancy after 30 weeks gestation
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 months
<b>Other Criteria</b>	PA applies to all. Product is approved for acute pain, defined as short-term pain not lasting longer than a 3-month period.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	



# DIFICID (fidaxomicin)

---

## Products Affected

- DIFICID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DIGOXIN IN OLDER PATIENTS

## Products Affected

- *digitek oral tablet 250 mcg*
- *digox oral tablet 250 mcg*
- *digoxin oral tablet 250 mcg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance) with result greater than or equal to 30 mL/min. Patient must have tried and failed to respond adequately to 0.125 mg of digoxin.
<b>Age Restrictions</b>	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
<b>Prescriber Restrictions</b>	PA not required for cardiology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all except cardiology. PA not required for doses less than or equal to 0.125 mg per day. Updated creatinine clearance since the previous authorization will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# DRONABINOL

## Products Affected

- *dronabinol*
- SYNDROS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use. If authorization is requested for treatment of nausea and vomiting associated with cancer therapy, documentation of previous conventional antiemetic therapies utilized is required.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. If the medication is being administered related to cancer treatment and is a full replacement for intravenous administration of antiemetic therapy within 48 hours of cancer treatment, it is covered as a Part B benefit. To be eligible for Part B coverage, the prescribing physician must indicate this on the prescription. If the medication is being requested for the use of anorexia associated with weight loss in patients with AIDS, approval may be covered under Part D.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# DROXIDOPA

---

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# DUOBRII (halobetasol/tazarotene)

---

## Products Affected

- DUOBRII

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential, documentation patient tried and failed augmented betamethasone dipropionate, clobetasol, fluocinonide 0.1%, halobetasol, or another Class I ultra-high potency topical steroid.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

## DUOPA (carbidopa/levodopa enteral suspension)

---

### Products Affected

- DUOPA ENTERAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Administration of non-selective monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation patient has a naso-jejunal tube for short-term administration or a PEG-J for long-term administration.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# DUPIXENT (dupilumab)

## Products Affected

- DUPIXENT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use. For atopic dermatitis, documentation of treatment with at least a moderate strength topical corticosteroid for at least four weeks, a contraindication to the use of topical corticosteroids, or therapy is not otherwise advisable. For moderate-to-severe asthma, either (1) submission of blood eosinophil count of at least 150 cells/mcL obtained within 6 weeks of therapy initiation or (2) documentation asthma requires daily oral corticosteroid for control. For chronic rhinosinusitis with nasal polyposis, documentation of treatment with an intranasal corticosteroid for at least three months, a contraindication to the use of intranasal corticosteroids, or therapy is not otherwise advisable.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to allergy, dermatology, immunology, otolaryngology/otorhinolaryngology, and pulmonology
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	PA applies to all. Continuation requires documentation of a positive response to therapy. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# EGRIFTA SV (tesamorelin)

---

## Products Affected

- EGRIFTA SV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, active malignancy, disruption of HPA axis due to hypophysectomy, hypopituitarism, pituitary tumor/surgery, head irradiation, or head trauma, use for weight loss
<b>Required Medical Information</b>	Diagnosis of covered use. Submission of pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Continuation of therapy requests require confirmation that the patient has demonstrated a clinical improvement (or maintenance of improvement once achieved) from baseline.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# EMFLAZA (deflazacort)

---

## Products Affected

- EMFLAZA

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, documentation of treatment failure with or intolerance to prednisone.
Age Restrictions	2 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# EMPAVELI (pegcetacoplan)

## Products Affected

- EMPAVELI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use confirmed by high-sensitivity flow cytometry, proof of vaccination against Streptococcus pneumoniae, Neisseria meningitidis, and Haemophilus influenzae type B or 2 weeks of antibacterial drug prophylaxis if the vaccines were administered within the last 2 weeks and therapy is required immediately, submission of lactate dehydrogenase level.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology, immunology, and nephrology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Because this medication is delivered subcutaneously through an infusion pump, it covered as a Part B benefit except for enrollees residing in a long-term care facility. PA applies to all when covered as a Part D benefit.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# EMSAM (selegiline transdermal)

---

## Products Affected

- EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	Pheochromocytoma, coadministration with carbamazepine or serotonergic drugs
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ENDOTHELIN RECEPTOR ANTAGONISTS

---

## Products Affected

- *ambrisentan*
- *bosentan*
- OPSUMIT
- TRACLEER ORAL TABLET SOLUBLE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy. For ambrisentan, idiopathic pulmonary fibrosis and moderate or severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential, submission of baseline AST, ALT, and bilirubin. For ambrisentan and Opsumit, submission of baseline hemoglobin level.
<b>Age Restrictions</b>	For ambrisentan and Opsumit, 18 years of age or older. For bosentan, 3 years of age or older.
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ENSPRYNG (satralizumab-mwge)

---

## Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Active hepatitis B infection, active or untreated latent tuberculosis (TB)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of confirmation patient has anti-aquaporin-4 (AQP4) antibody-positive NMOSD, submission of baseline latent TB screening test (Mantoux tuberculin skin test [a.k.a. PPD test] or interferon-gamma release assay [IGRA]), attestation patient does not have any active infection.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and ophthalmology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# EPIDIOLEX (cannabidiol)

---

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	1 year of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ERIVEDGE (vismodegib)

---

## Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ERLOTINIB

---

## Products Affected

- *erlotinib hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For non-small cell lung cancer, submission of FDA-approved test confirming presence of EGFR exon 19 deletion or exon 21 L858R substitution mutation and prior treatments used.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# ESBRIET (pirfenidone)

---

## Products Affected

- ESBRIET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	End-stage renal disease on dialysis, severe (Child-Pugh class C) hepatic impairment
<b>Required Medical Information</b>	Diagnosis of covered use, submission of liver function tests or Child-Pugh status.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated liver function testing or Child-Pugh score since the previous authorization will be required for subsequent reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# EVEROLIMUS

---

## Products Affected

- AFINITOR DISPERZ
- AFINITOR ORAL TABLET 10 MG
- *everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with dual strong CYP3A4/P-glycoprotein inhibitors
Required Medical Information	Diagnosis of covered use and submission of pregnancy status for female patients of childbearing potential. For renal cell carcinoma, documented prior use of sunitinib or sorafenib. For postmenopausal women with advanced hormone receptor-positive, HER-2 negative breast cancer, documentation of treatment failure with letrozole or anastrozole and confirmation drug is being used in combination with exemestane.
Age Restrictions	1 year of age or older
Prescriber Restrictions	Restricted to neurology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# FARYDAK (panobinostat)

## Products Affected

- FARYDAK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, recent myocardial infarction or unstable angina, coadministration with strong CYP3A4 inducers, antiarrhythmic drugs, or drugs that prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, documentation that the patient has received at least 2 prior regimens including bortezomib and an immunomodulatory agent, submission of baseline ECG documenting QTcF is less than 450 msec prior to initiation, submission of baseline serum electrolytes including potassium and magnesium, submission of baseline CBC documenting platelet count is at least $100 \times 10^9/L$ and absolute neutrophil count is at least $1.5 \times 10^9/L$ , submission of baseline liver function tests including AST, ALT, and total bilirubin. For multiple myeloma, confirmation drug will be given with dexamethasone and bortezomib.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	Initially 8 cycles. An additional 8 cycles if clinical benefit seen.
<b>Other Criteria</b>	PA applies to new starts only. An additional 8 cycles may be approved only if there is an absence of unresolved severe or medically significant toxicity.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# FENTANYL TRANSMUCOSAL

---

## Products Affected

- *fentanyl citrate buccal*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients not tolerant to the effects of a chronic opioid, treatment of acute or postoperative pain including headache, migraines, or dental pain
<b>Required Medical Information</b>	Diagnosis of covered use, verified claim or documentation of patient's morphine milligram equivalent opioid dose.
<b>Age Restrictions</b>	For the buccal tablet, 18 years of age or older. For the lozenge, 16 years of age or older.
<b>Prescriber Restrictions</b>	PA not required for oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all except oncology.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# FINTEPLA (fenfluramine)

---

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Hepatic impairment, moderate or severe renal impairment, administration of monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance) and liver function testing or Child-Pugh score.
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# FIRDAPSE (amifampridine)

---

## Products Affected

- FIRDAPSE

PA Criteria	Criteria Details
Exclusion Criteria	History of seizure
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# FIRST-GENERATION ANTIHISTAMINES IN OLDER PATIENTS

## Products Affected

- *carbinoxamine maleate oral solution*
- *carbinoxamine maleate oral tablet 4 mg*
- *clemastine fumarate oral tablet 2.68 mg*
- *cyproheptadine hcl oral*
- *diphenhydramine hcl oral elixir*
- *hydroxyzine hcl oral tablet*
- *hydroxyzine pamoate oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use. For carbinoxamine or cyproheptadine for dermatographism, documentation patient tried and had an inadequate response to a second-generation antihistamine. For hydroxyzine for pruritus, documentation patient tried and had an inadequate response to a second-generation antihistamine. For hydroxyzine for anxiety, documentation patient has tried and had an inadequate response to at least 2 other FDA-approved products for the management of anxiety OR documentation medication is being used as a sedative before and after general anesthesia.
<b>Age Restrictions</b>	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. First-generation antihistamines are anticholinergic medications considered high-risk in older patients due to risks of confusion, dry mouth, constipation, and decreased clearance with advanced age.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# FOTIVDA (tivozanib)

---

## Products Affected

- FOTIVDA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncontrolled hypertension, severe hepatic impairment, coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of previous systemic therapies used to treat renal cell carcinoma including the failure of at least one prior VEGFR inhibitor, pregnancy status for female patients of childbearing potential, confirmation patient has not had episodes of symptomatic heart failure or unstable angina, a myocardial infarction, an arterial thrombotic event, or a significant bleeding event in the 6 months preceding the prior authorization request.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# FUMARATES FOR MULTIPLE SCLEROSIS

---

## Products Affected

- BAFIERTAM
- VUMERITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Hypersensitivity to dimethyl fumarate, coadministration with another fumarate. For Vumerity, moderate or severe renal impairment.
<b>Required Medical Information</b>	Diagnosis of covered use. For Vumerity, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. For approval of Bafiertam or Vumerity, the patient must have tried and failed to have an adequate response to or had an intolerance to dimethyl fumarate. Updated creatinine clearance since the previous authorization will be required for subsequent annual reauthorizations of Vumerity.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# GALAFOLD (migalastat)

---

## Products Affected

- GALAFOLD

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe renal impairment (eGFR less than 30 mL/min/1.73 m <sup>2</sup> ) or end-stage renal disease requiring dialysis
<b>Required Medical Information</b>	Diagnosis of covered use, documentation that the patient has an amenable galactosidase alpha gene variant (see section 12.1, table 2 of package insert for full list) based on in vitro assay data as interpreted by a clinical genetics professional.
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# GATTEX (teduglutide)

## Products Affected

- GATTEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline laboratory values including bilirubin, alkaline phosphatase, lipase, and amylase obtained within 6 months prior to starting therapy. For adults 18 years of age or older only, submission of documentation that a colonoscopy (or alternate imaging) of the entire colon with polyp removal was performed within 6 months prior to starting treatment.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. For adults 18 years of age or older, continuation of therapy requires submission of findings from a follow-up colonoscopy or alternate imaging result at the end of 1 year of teduglutide treatment. Subsequent imaging should be performed every 5 years, or sooner if polyps are found at the 1-year mark.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# GAVRETO (pralsetinib)

---

## Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inhibitors
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of RET gene fusion or mutation, attestation patient does not have uncontrolled hypertension, pregnancy status for female patients of childbearing potential.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# GILOTRIF (afatinib)

---

## Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For non-small cell lung cancer, submission of positive FDA-approved test for non-resistant epidermal growth factor receptor mutations. For metastatic, squamous non-small cell lung cancer, documentation of progression after platinum-based chemotherapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# GnRH ANTAGONISTS

## Products Affected

- ELIGARD
- FIRMAGON (240 MG DOSE)
- FIRMAGON SUBCUTANEOUS SOLUTION RECONSTITUTED 80 MG
- *leuprolide acetate injection*
- LUPANETA PACK
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)
- LUPRON DEPOT-PED (1-MONTH) INTRAMUSCULAR KIT 11.25 MG, 15 MG
- LUPRON DEPOT-PED (3-MONTH) INTRAMUSCULAR KIT 30 MG (PED)
- TRELSTAR MIXJECT

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For prostate cancer, documentation of baseline prostate-specific antigen and serum testosterone level.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology, oncology, endocrinology, gynecology, and urology
Coverage Duration	For endometriosis and uterine fibroids, 6 months. For all other indications, 1 year.
Other Criteria	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# GROWTH HORMONE

## Products Affected

- GENOTROPIN
- GENOTROPIN MINIQUICK
- HUMATROPE
- NORDITROPIN FLEXPRO SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 10 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 20 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- NUTROPIN AQ NUSPIN 5 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of growth failure, submission of IGF-1 levels, height, weight, creatinine clearance (or serum creatinine), fasting blood glucose, and bone age if applicable based on patient age and diagnosis.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to endocrinology and nephrology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated IGF-1 level, bone age (if applicable based on patient age and diagnosis) height, weight, creatinine clearance (or serum creatinine), and fasting glucose since the previous authorization will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# HEREDITARY ANGIOEDEMA THERAPIES, ACUTE

## Products Affected

- *icatibant acetate*
- RUCONEST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for prophylactic hereditary angioedema therapy. For Ruconest, acute laryngeal attacks.
<b>Required Medical Information</b>	Diagnosis of covered use. For Ruconest, documentation of the patient's typical attack presentation/symptoms.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to allergy, dermatology, hematology, or immunology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# HEREDITARY ANGIOEDEMA THERAPIES, MAINTENANCE

## Products Affected

- HAEGARDA
- ORLADEYO
- *takhzyro*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Requests for acute hereditary angioedema therapy (attacks). For Orladeyo, end-stage renal disease.
<b>Required Medical Information</b>	Diagnosis of covered use, submission of objective or subjective documentation that prophylactic therapy is medically necessary, including, but not limited to activity of disease and disease burden, the frequency of HAE attacks, and quality of life.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to allergy, dermatology, hematology, or immunology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval of either Haegarda or Orladeyo for patients 12 years of age and older, the patient must have tried and failed to have an adequate response to or had an intolerance or contraindication to Takhzyro. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# HETLIOZ (tasimelteon)

---

## Products Affected

- HETLIOZ

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with strong CYP1A2 inhibitors or CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use. For Smith-Magenis Syndrome patients only, documentation of genetic testing results confirming diagnosis is required.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to neurology and sleep medicine
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. For non-24-hour sleep-wake disorder, patients are required to be totally blind to match the population in which tasimelteon was studied.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# HIGH-RISK DRUGS IN OLDER PATIENTS

---

## Products Affected

- BUTISOL SODIUM ORAL TABLET 30 MG
- CHLORPROPAMIDE ORAL TABLET 100 MG
- *meprobamate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	All 3 of the following criteria must be met: (1) diagnosis of covered use, (2) documentation provider is aware medication is considered a high-risk medication for elderly patients according to the Centers for Medicare and Medicaid Services, and (3) documentation that the benefits of the drug outweigh the potential risks to the patient.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	

# IBRANCE (palbociclib)

---

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of genetic tumor testing showing that the primary tumor type is HR-positive, HER2-negative, confirmation that the treatment regimen will include concomitant use of an aromatase inhibitor or fulvestrant, submission of baseline CBC.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ICLUSIG (ponatinib)

---

## Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	Newly diagnosed chronic phase CML
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# IDHIFA (enasidenib)

---

## Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of IDH2 mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ILARIS

## Products Affected

- ILARIS SUBCUTANEOUS SOLUTION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Positive TB test, coadministration with TNF inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of TB skin test result obtained within past 12 months, documentation that patient has received all recommended vaccinations as appropriate including pneumococcal vaccine and inactivated influenza vaccine prior to initiation of therapy. For CAPS, confirmed diagnosis including genetic testing for variant FCAS or MWS and documentation patient is not receiving concomitant TNF inhibitor therapy. For SJIA, submission of CBC including platelet count and confirmed diagnosis defined by prominence of systemic and inflammatory features including spiking fevers, rash, swelling and inflammation of lymph nodes, liver, and spleen, and high white blood cell and platelet counts.
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	For SJIA, restricted to rheumatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated TB skin test result obtained within the past 12 months and objective documentation of positive patient response or maintenance of response will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# IMBRUVICA (ibrutinib)

---

## Products Affected

- IMBRUVICA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For chronic graft versus host disease, documentation of treatment failure with any other systemic immunosuppressive agent.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology, oncology, and transplant specialty
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# IMCIVREE (setmelanotide)

## Products Affected

- IMCIVREE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe renal impairment, end-stage renal disease
<b>Required Medical Information</b>	Diagnosis of covered use, including submission of genetic testing showing homozygous or compound heterozygous gene variants in POMC, PCSK1, or LEPR genes interpreted as pathogenic, likely pathogenic, or of uncertain clinical significance and body mass index (BMI) greater than 30 kg/m <sup>2</sup> in adults or greater than the 97th percentile in children.
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Initially 16 weeks, then 1 year
<b>Other Criteria</b>	PA applies to all. For re-authorization at the 16-week point, submission of clinical documentation attesting to at least 5% weight loss from baseline (or at least 5% BMI from baseline in patients with continued growth potential) is required. Not FDA-approved for other types or causes of obesity, and therefore requests for these uses will not be approved.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

# IMMUNE GLOBULIN

## Products Affected

- BIVIGAM INTRAVENOUS SOLUTION 5 GM/50ML GM/100ML, 10 GM/200ML, 20 GM/200ML, 20 GM/400ML, 5 GM/100ML, 5 GM/50ML
- CARIMUNE NF INTRAVENOUS SOLUTION RECONSTITUTED 12 GM, 6 GM
- FLEBOGAMMA DIF
- GAMMAGARD
- GAMMAGARD S/D LESS IGA
- GAMMAKED INJECTION SOLUTION 1 GM/10ML, 10 GM/100ML, 20 GM/200ML, 5 GM/50ML
- GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 2 GM/20ML, 2.5 GM/50ML, 20 GM/200ML, 25 GM/500ML, 5 GM/100ML, 5 GM/50ML
- GAMUNEX-C
- OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 10 GM/100ML, 10 GM/200ML, 2 GM/20ML, 2.5 GM/50ML, 20 GM/200ML, 25 GM/500ML, 5 GM/100ML, 5 GM/50ML
- PRIVIGEN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	IgA-deficient patients with antibodies against IgA and a history of hypersensitivity. For IM forms, severe thrombocytopenia or coagulation disorder that would contraindicate an IM injection.
<b>Required Medical Information</b>	Diagnosis of covered use. For ITP, submission of platelet count. For CLL, IgG level less than 600 mg/dL and recent history of serious bacterial infection requiring either oral or IV antibiotic therapy. For CIDP, unequivocal diagnosis and documentation patient is refractory or intolerant to prednisone or azathioprine given in therapeutic doses over at least 3 months. For passive immunization against varicella, confirmation that the patient is immunosuppressed and cannot receive varicella-zoster immune globulin.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	For acute conditions/new starts, 3 months. For renewal of chronic conditions, 1 year.
<b>Other Criteria</b>	PA applies to all. For continuation of any diagnosis, documentation of the clinical response to therapy must be submitted. For IV formulations, covered as a Part B benefit if administered in the home for the treatment of primary immune deficiency. For any other combination of treatment site and indication, additional information may need to be submitted to determine if the immune globulin will be covered as a Part B or Part D benefit.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INJECTABLE TESTOSTERONE

---

## Products Affected

- *testosterone cypionate injection solution 200 mg/ml*
- *testosterone cypionate intramuscular solution 100 mg/ml, 200 mg/ml*
- *testosterone enanthate intramuscular solution*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum testosterone level, documentation that patient has been evaluated for the presence of breast and prostate cancer prior to initiation of therapy.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PA not required for urology or endocrinology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all except when prescribed by urology or endocrinology. Documentation of clinically relevant response to therapy (including, but not limited to submission of updated serum testosterone level) will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INLYTA (axitinib)

---

## Products Affected

- INLYTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), uncontrolled hypertension, evidence of untreated brain metastasis, recent active gastrointestinal bleeding, coadministration with strong CYP3A4/5 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of laboratory values including baseline ALT, AST, bilirubin, submission of baseline blood pressure reading. If axitinib is being used as a single agent, submission of prior therapy or therapies tried and failed.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INQOVI (decitabine/cedazuridine)

---

## Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of complete blood count, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to new starts only. Continuation of initial therapy beyond 6 months requires (a) confirmation of no disease progression and (b) attestation the patient is having no serious adverse events from treatment.
Indications	All Medically-accepted Indications.
Off Label Uses	

# INREBIC (fedratinib)

---

## Products Affected

- INREBIC

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with moderate or strong CYP3A4 inducers or dual CYP3A4/CYP2C19 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of thiamine level and platelet count.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval, the patient must have tried and failed to have an adequate response to or had an intolerance to ruxolitinib.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INTRANASAL SEIZURE MEDICATIONS

---

**Products Affected**

- NAYZILAM
- VALTOCO 10 MG DOSE
- VALTOCO 15 MG DOSE
- VALTOCO 20 MG DOSE
- VALTOCO 5 MG DOSE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute narrow-angle glaucoma
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INTRON A (interferon alfa-2b)

## Products Affected

- INTRON A

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Autoimmune hepatitis, decompensated liver disease
<b>Required Medical Information</b>	Diagnosis of covered use, submission of triglyceride levels, hemoglobin, complete and differential white blood cell counts, platelet count, serum electrolytes, ALT, serum bilirubin level, serum albumin level, and TSH. For malignant melanoma, submission of the date of surgical treatment. For AIDS-related Kaposi's sarcoma, submission of total CD4 count. For chronic hepatitis C, submission of HCV RNA, prothrombin time, baseline serum creatinine level, and laboratory confirmation of hepatitis C virus, and documentation of previous response to therapy if applicable. For chronic hepatitis B, submission of prothrombin time and documentation patient has been serum HBsAG positive for at least 6 months with evidence of HBV replication.
<b>Age Restrictions</b>	For hairy cell leukemia, malignant melanoma, follicular lymphoma, condylomata acuminata, or AIDS-related Kaposi's sarcoma, 18 years of age or older. For chronic hepatitis C, 3 years of age or older. For chronic hepatitis B, 1 year of age or older.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Depends on covered use. See "Other Criteria" section.
<b>Other Criteria</b>	PA applies to new starts only. For hairy cell leukemia, the coverage duration is 6 months. For condylomata acuminata, 3 weeks per course, and at least 12 weeks must pass in between multiple courses in order to be reauthorized. For Kaposi's sarcoma, 16 weeks. For hepatitis B infection, 24 weeks. For all other indications/uses, 1 year. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# INVEGA HAFYERA (paliperidone 6-month injectable suspension)

## Products Affected

- INVEGA HAFYERA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension or at least one 3-month injection of 3-month paliperidone palmitate extended-release injectable suspension.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# INVEGA TRINZA (paliperidone 3-month injectable suspension)

## Products Affected

- INVEGA TRINZA INTRAMUSCULAR SUSPENSION  
PREFILLED SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of at least 4 months' treatment with 1-month paliperidone palmitate extended-release injectable suspension.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# IRESSA (gefitinib)

---

## Products Affected

- IRESSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation of EGFR exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ISTURISA (osilodrostat)

---

## Products Affected

- ISTURISA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, attestation pituitary gland surgery is not an option for the patient or has not been curative, submission of baseline serum potassium and magnesium levels.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to all. Continuation requires documentation of clinically relevant response to therapy, including, but not limited to urine free cortisol level.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ITRACONAZOLE

---

## Products Affected

- *itraconazole oral*
- TOLSURA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, fungal culture result identifying causative organism or positive KOH result.
Age Restrictions	
Prescriber Restrictions	PA not required for infectious diseases
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# JAKAFI (ruxolitinib)

---

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For myelofibrosis, a platelet count less than $50 \times 10^9/L$ with either concomitant estimated creatinine clearance between 15 and 59 mL/min, end-stage renal disease not on dialysis, or any degree of hepatic impairment
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline platelet count, ALT, AST, and bilirubin, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance). For polycythemia vera, documented intolerance or inadequate response to hydroxyurea. For acute graft-versus-host disease, documented inadequate response to systemic corticosteroids.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology, oncology, and transplant specialty
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# JUXTAPID (lomitapide)

## Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, moderate or severe hepatic impairment (Child-Pugh class B or C), active liver disease, coadministration with moderate or strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline lab values including ALT, AST, alkaline phosphatase, total bilirubin, baseline LDL-C, total cholesterol (TC), apoB, and non-HDL-C, pregnancy status for female patients of childbearing potential, submission of renal indices, documentation of contraindication to or treatment failure with evolocumab.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology, lipidology, and endocrinology with experience in and a focus on lipid management
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	PA applies to all. Submission of LDL level drawn after the initial LDL level submission documenting clinically significant response to therapy will be required for reauthorization. For approval, the patient must have tried and failed to have an adequate response to, had an intolerance to, or have a contraindication to therapy with evolocumab.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# JYNARQUE (tolvaptan)

---

## Products Affected

- JYNARQUE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of signs or symptoms of significant liver impairment or injury (not including uncomplicated polycystic liver disease), uncorrected abnormal blood sodium concentrations, inability to sense or respond to thirst, hypovolemia, uncorrected urinary outflow obstruction, anuria, coadministration with strong CYP3A inhibitors or inducers or desmopressin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum sodium concentration.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to nephrology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# KALYDECO (ivacaftor)

---

## Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of cystic fibrosis mutation test result and baseline ALT and AST.
Age Restrictions	
Prescriber Restrictions	Restricted to pulmonology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# KERENDIA (finerenone)

---

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Adrenal insufficiency, estimated glomerular filtration rate (eGFR) less than 25 mL/min/1.73 m <sup>2</sup> , severe (Child-Pugh class C) hepatic impairment, coadministration with strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of estimated glomerular filtration rate (eGFR) and baseline serum potassium level.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. For approval, the patient must have documentation of a trial of Farxiga.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# KETOCONAZOLE ORAL

---

## Products Affected

- *ketoconazole oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Acute or chronic liver disease, treatment of fungal meningitis or fungal infections of the skin or nails
<b>Required Medical Information</b>	Ketoconazole is being requested for the treatment of culture-proven systemic blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis, submission of baseline ALT, AST, total bilirubin, alkaline phosphatase, prothrombin time and INR, confirmation from the prescriber that the potential benefits of therapy outweigh the risks.
<b>Age Restrictions</b>	2 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 months
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# KEVEYIS (dichlorphenamide)

---

## Products Affected

- KEVEYIS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use of high dose aspirin, severe pulmonary disease limiting compensation to metabolic acidosis, hepatic insufficiency
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum potassium.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Initially 2 months, then 1 year
<b>Other Criteria</b>	PA applies to all. Documentation of a positive response to therapy will be required for initial reauthorization after the first 2 months. Maintenance of a clinical benefit will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# KORLYM (mifepristone)

---

**Products Affected**

- KORLYM

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment, uncorrected hypokalemia, female patients with a history of unexplained vaginal bleeding, endometrial hyperplasia with atypia, or endometrial carcinoma, patients using systemic corticosteroids for life-saving purposes, coadministration with strong CYP3A4 inducers, simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges
<b>Required Medical Information</b>	Diagnosis of covered use, attestation surgery is not an option for the patient or has not been curative, submission of baseline serum potassium, AST, ALT, and alkaline phosphatase, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# KOSELUGO (selumetinib)

---

## Products Affected

- KOSELUGO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of left ventricular ejection fraction, Child-Pugh score or liver function testing results, and pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	Initiation: 2-17 years of age. Continuation: 2 years of age or older.
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	PA applies to new starts only. Selumetinib is indicated in pediatric patients and will not be approved for adults unless the patient started on the medication prior to 18 years of age. Continuation of initial therapy beyond 6 months requires (a) documentation of any positive clinical response and (b) attestation the patient is having no serious adverse events to treatment.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# LAPATINIB

---

## Products Affected

- *lapatinib ditosylate*

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia, uncorrected hypomagnesemia
Required Medical Information	Diagnosis of covered use, submission of baseline left ventricular ejection fraction, potassium and magnesium levels, pregnancy status for female patients of childbearing potential, and depending on indication, confirmation that the treatment regimen will include concomitant use of either capecitabine or letrozole. For patients who will be using lapatinib with capecitabine, submission of prior therapies tried and failed.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# LENVIMA (lenvatinib)

---

## Products Affected

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected electrolyte abnormalities, uncontrolled hypertension
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline blood pressure, comprehensive metabolic panel, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# LEUKINE (sargramostim, GM-CSF)

---

## Products Affected

- LEUKINE INJECTION SOLUTION RECONSTITUTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of WBC count and ANC.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# LIDOCAINE TRANSDERMAL PATCHES

---

## Products Affected

- *lidocaine external patch 5 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. FDA-approved only for postherpetic neuralgia. Requests for other indications will not be approved.
Indications	All FDA-approved Indications.
Off Label Uses	

# LONSURF (trifluridine/tipiracil)

---

## Products Affected

- LONSURF

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe hepatic impairment, febrile neutropenia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of prior therapies used, submission of baseline CBC, absolute neutrophil count, ALT, AST, and bilirubin, pregnancy status for female patients of childbearing potential, documentation of KRAS status.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# LORBRENA (lorlatinib)

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers, uncontrolled hypertension
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of ALK-positive tumor, baseline blood pressure, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# LUCEMYRA (lofexidine)

---

## Products Affected

- LUCEMYRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Congenital long QT syndrome, severe coronary insufficiency, recent myocardial infarction, cerebrovascular disease, chronic renal failure
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient will be using medication in conjunction with a comprehensive management program for the treatment of opioid use disorder.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	14 days
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# LUMAKRAS (sotorasib)

---

## Products Affected

- LUMAKRAS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers, coadministration with proton pump inhibitors or H2 receptor antagonists
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test result confirming presence of KRAS G12C mutations, submission of previous systemic treatment(s) tried.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# LUPKYNIS (voclosporin)

## Products Affected

- LUPKYNIS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with strong CYP3A4 inhibitors, moderate or strong CYP3A4 inducers, or cyclophosphamide, hypertensive emergency or a baseline blood pressure above 165/105 mmHg
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient will be taking concurrently with mycophenolate mofetil and corticosteroids, submission of estimated glomerular filtration rate (eGFR), baseline blood pressure, pregnancy status for female patients of childbearing potential. If the patient's eGFR is less than or equal to 45 mL/min/1.73 m <sup>2</sup> , attestation that prescriber believes benefits of therapy outweigh the potential risks to the patient.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Initially 6 months, then 1 year
<b>Other Criteria</b>	PA applies to all. For approval, the patient must have tried and failed to have an adequate response to or had an intolerance/contraindication to Benlysta (belimumab). Documentation of a positive response to therapy will be required for initial reauthorization after the first 6 months. Maintenance of a clinical benefit, attestation that prescriber believes benefits of continuing therapy outweigh the potential risks to the patient, and updated estimated glomerular filtration rate (eGFR) and blood pressure since the previous authorization will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# LYNPARZA (olaparib)

---

## Products Affected

- LYNPARZA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of applicable mutations and previous therapies tried and failed depending on cancer type as necessary, submission of baseline CBC.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and urology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# MAVENCLAD (cladribine)

## Products Affected

- MAVENCLAD (10 TABS)
- MAVENCLAD (4 TABS)
- MAVENCLAD (5 TABS)
- MAVENCLAD (6 TABS)
- MAVENCLAD (7 TABS)
- MAVENCLAD (8 TABS)
- MAVENCLAD (9 TABS)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Current malignancy, pregnancy, HIV or other active chronic infection (e.g., hepatitis or tuberculosis), lymphocyte count below normal limit before first course or less than 800 cells/microliter before second course, creatinine clearance below 60 mL/min, Child-Pugh score greater than 6
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential, submission of lymphocyte count, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Documentation of a positive response to therapy, confirmation the patient has no active infection, and updated lymphocyte count and creatinine clearance since the previous authorization will be required for reauthorization. After the completion of 2 treatment courses (2 years' treatment), additional treatment courses are not recommended over the following 2 years because of malignancy risk. Re-initiating treatment after those 2 years have passed has not been studied. Requests for therapy for a combined total of greater than 2 years will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# MAVYRET (glecaprevir/pibrentasvir)

---

## Products Affected

- MAVYRET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe hepatic impairment (Child-Pugh class B or C), coadministration with rifampin or atazanavir
<b>Required Medical Information</b>	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV), documentation of whether cirrhosis is present or not and whether or not it is compensated or decompensated, confirmation a test for HBV infection (HBsAg and anti-HBc) was completed.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Criteria for coverage duration will be applied consistent with current AASLD-IDSA guidance.
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# MECASERMIN

---

## Products Affected

- INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Patients with closed epiphyses
Required Medical Information	Diagnosis of covered use, documentation of primary insulin-like growth factor (IGF-1) deficiency or growth hormone gene deletion in patients who have developed neutralizing antibodies to growth hormone, submission of IGF-1 level and growth hormone level.
Age Restrictions	
Prescriber Restrictions	Restricted to endocrinology and nephrology
Coverage Duration	6 months
Other Criteria	PA applies to all. Updated IGF-1 and growth hormone levels since the previous authorization will be required for subsequent reauthorizations. Mecasermin is not indicated as a growth hormone replacement and will not be approved for this indication.
Indications	All Medically-accepted Indications.
Off Label Uses	

# MEGESTROL IN OLDER PATIENTS

---

## Products Affected

- *megestrol acetate oral suspension 40 mg/ml, 400 mg/10ml, 625 mg/5ml*

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
Prescriber Restrictions	PA not required for hematology or oncology
Coverage Duration	1 year
Other Criteria	PA applies to all except hematology and oncology.
Indications	All Medically-accepted Indications.
Off Label Uses	

# MEKINIST (trametinib)

---

## Products Affected

- MEKINIST

PA Criteria	Criteria Details
Exclusion Criteria	Progression of disease on prior BRAF-inhibitor therapy
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of BRAF V600E or V600K mutation. For non-small cell lung cancer and thyroid cancer, attestation that therapy will be used in combination with dabrafenib.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# METHAMPHETAMINE

---

## Products Affected

- *methamphetamine hcl*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Use for exogenous obesity, patients with glaucoma, advanced arteriosclerosis, symptomatic cardiovascular disease, moderate to severe hypertension, hyperthyroidism, or a history of drug abuse, administration of non-selective monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use. For patients 65 years of age and older, attestation provider is aware medication is considered a high-risk medication for elderly patients according to the Centers for Medicare and Medicaid Services (CMS) and that the benefits of methamphetamine therapy outweigh the potential risks to the patient.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. PA will not be authorized if using for exogenous obesity (excluded category per CMS).
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# METHOTREXATE INJECTABLE (SUBCUTANEOUS)

---

## Products Affected

- OTREXUP SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.4ML, 12.5 MG/0.4ML, 15 MG/0.4ML, 17.5 MG/0.4ML, 20 MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML
- RASUVO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML

- REDITREX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	Pregnancy
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of intolerance or inadequate response to oral or non-subcutaneous injectable forms of methotrexate.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to rheumatology and dermatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# METHYLTESTOSTERONE

---

## Products Affected

- METHITEST
- *methyltestosterone oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Male patients with breast or prostate cancer, women who are or may become pregnant
<b>Required Medical Information</b>	Diagnosis of covered use. For patients 65 years of age and older, attestation provider is aware medication is considered a high-risk medication for elderly patients according to the Centers for Medicare and Medicaid Services (CMS) and that the benefits of methyltestosterone therapy outweigh the potential risks to the patient.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# MIGLUSTAT

---

## Products Affected

- *miglustat*

PA Criteria	Criteria Details
Exclusion Criteria	Severe renal impairment (CrCl less than 30 mL/min)
Required Medical Information	Diagnosis of covered use, documentation that enzyme replacement is not a therapeutic option.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# MYALEPT (metreleptin)

---

## Products Affected

- MYALEPT

PA Criteria	Criteria Details
Exclusion Criteria	General obesity not associated with congenital leptin deficiency
Required Medical Information	Diagnosis of covered use, submission of leptin level laboratory test result confirming leptin deficiency, baseline HbA1c, fasting glucose, fasting triglyceride levels, and weight.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated patient weight, HbA1c, fasting glucose, and fasting triglyceride levels since the previous authorization will be required for subsequent annual reauthorizations. Metreleptin is not established as a treatment for nonalcoholic steatohepatitis, complications of partial lipodystrophy, HIV-related lipodystrophy, or metabolic disease without generalized lipodystrophy, and submissions for these uses will not be approved.
Indications	All Medically-accepted Indications.
Off Label Uses	

# MYCAPSSA (octreotide)

---

## Products Affected

- MYCAPSSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of prior use of either injectable octreotide or lanreotide and attestation to its successful treatment of acromegaly using clinical biomarkers or chart notes.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# MYFEMBREE (relugolix/estradiol/norethindrone)

## Products Affected

- MYFEMBREE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known liver impairment or disease, known osteoporosis, undiagnosed abnormal uterine bleeding, women at increased risk of or current/a history of thrombotic or thromboembolic disorders (including women over 35 years of age who smoke and women with uncontrolled hypertension), current/history of breast cancer or other hormone-sensitive cancer
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is premenopausal, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# MYTESI (crofelemer)

---

## Products Affected

- MYTESI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, attestation infectious causes of diarrhea have been ruled out.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# NAMZARIC (memantine and donepezil)

---

## Products Affected

- NAMZARIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of documentation that the patient has been stabilized on donepezil 10 mg daily.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# NATPARA (parathyroid hormone)

---

## Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation that (albumin-corrected) serum calcium is greater than 7.5 mg/dL and confirmation that 25-hydroxyvitamin D stores are sufficient.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	



# NERLYNX (neratinib)

---

## Products Affected

- NERLYNX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with proton pump inhibitors, strong CYP3A4 inhibitors, moderate CYP3A4 and P-glycoprotein dual inhibitors, or moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing showing that the primary tumor type is HER2-positive, confirmation member has completed adjuvant trastuzumab-based therapy or will be using in combination with capecitabine, submission of baseline liver function tests, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# NEXAVAR (sorafenib)

---

## Products Affected

- NEXAVAR

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# NINLARO (ixazomib)

## Products Affected

- NINLARO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, documentation that medication will be administered concomitantly with lenalidomide and dexamethasone, documentation of prior therapy regimen for multiple myeloma, submission of baseline platelet count, absolute neutrophil count, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	Initially 3 months, then 1 year
<b>Other Criteria</b>	PA applies to new starts only. For continuation, documentation of platelet count greater than 30,000/mm <sup>3</sup> , ANC greater than 500/mm <sup>3</sup> , and Grade 1 or lower non-hematological toxicities (including rash, peripheral neuropathies) required.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# NITISINONE

---

## Products Affected

- *nitisinone*
- NITYR
- ORFADIN ORAL CAPSULE 20 MG
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to all. Updated liver function tests, urine succinylacetone levels, alpha-fetoprotein level, serum tyrosine level, and serum phenylalanine level since the previous authorization will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# NUEDEXTA (dextromethorphan and quinidine)

---

## Products Affected

- NUEDEXTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Prolonged QT interval, congenital long QT syndrome, heart failure, history suggestive of torsades de pointes, AV block without implanted pacemaker, administration of non-selective monoamine oxidase inhibitors within 14 days of initiation
<b>Required Medical Information</b>	Diagnosis of covered use, submission of ECG (specifically QT interval).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. The medication will not be approved for agitation or Alzheimer's disease without pseudobulbar affect as this is considered an off-label use.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# NUPLAZID (pimavanserin)

---

## Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis unrelated to Parkinson's disease psychosis, cardiac arrhythmias, symptomatic bradycardia, congenital QT prolongation, coadministration with moderate or strong CYP3A4 inducers, drugs that prolong the QT interval, hypokalemia, hypomagnesemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of comprehensive metabolic panel and ECG (specifically QT interval).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# OCALIVA (obeticholic acid)

## Products Affected

- OCALIVA ORAL TABLET 10 MG, 5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Complete biliary obstruction, decompensated cirrhosis (Child-Pugh B or C) or prior decompensation event, compensated cirrhosis with evidence of portal hypertension (e.g., ascites, gastroesophageal varices, persistent thrombocytopenia)
<b>Required Medical Information</b>	Diagnosis of covered use, documentation either (1) drug will be used in combination with ursodeoxycholic acid (UDCA) and UDCA has been used for 1 year or (2) patient had intolerance to UDCA, submission of baseline LFTs including ALP and total bilirubin, attestation patient does not have evidence of portal hypertension and has not had a prior decompensation event.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	Initially 3 months, then 1 year
<b>Other Criteria</b>	PA applies to all. Updated ALP obtained within the previous 3 months will be required for subsequent authorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ODOMZO (sonidegib)

---

## Products Affected

- ODOMZO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, coadministration with strong CYP3A4 inhibitors or moderate or strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is not a candidate for surgery or radiation therapy or carcinoma has recurred following surgery or radiation therapy, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# OFEV (nintedanib)

---

## Products Affected

- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe (Child-Pugh class B or C) hepatic impairment, coadministration of a dual P-glycoprotein/CYP3A4 inducer
<b>Required Medical Information</b>	Diagnosis of covered use, submission of liver function tests or Child-Pugh status, pregnancy status in female patients of childbearing potential. For chronic fibrosing interstitial lung diseases with a progressive phenotype and systemic sclerosis-associated interstitial lung disease diagnoses, submission of HRCT scan showing fibrosis affecting at least 10% of the lungs within the previous 12 months.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to pulmonology or rheumatology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated liver function testing or Child-Pugh score since the previous authorization will be required for subsequent reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ONUREG (azacitidine)

---

## Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of absolute neutrophil count, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to new starts only. Attestation of clinical benefit or stabilization and absence of unacceptable toxicity will be required for reauthorization. This dosage form is not intended to be a substitute for or substituted for injectable azacitidine.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ORENITRAM (treprostinil)

---

## Products Affected

- ORENITRAM

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe (Child-Pugh class B or C) hepatic impairment
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ORGOVYX (relugolix)

---

## Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and urology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ORIAHNN (elagolix/estradiol/norethindrone)

## Products Affected

- ORIAHNN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, known liver impairment or disease, known osteoporosis, undiagnosed abnormal uterine bleeding, women at increased risk of or a history of thrombotic or thromboembolic disorders (including women over 35 years of age who smoke and women with uncontrolled hypertension), current/history of breast cancer or other hormone-sensitive cancer, coadministration with OATP1B1 inhibitors or strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is premenopausal, submission of baseline blood pressure, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ORILISSA (elagolix)

## Products Affected

- ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, severe hepatic impairment (Child-Pugh class C), known osteoporosis, coadministration with OATP1B1 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology and gynecology
<b>Coverage Duration</b>	Up to 24 months based on liver function and coexisting dyspareunia. See "Other Criteria" section.
<b>Other Criteria</b>	PA applies to all. For endometriosis with dyspareunia or in women with moderate hepatic impairment, 6 months. For endometriosis without dyspareunia, 150 mg daily for 24 months. Use of this drug for more than 2 years increases risk of bone loss and requests for therapy for more than 2 years will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ORKAMBI (lumacaftor/ivacaftor)

---

## Products Affected

- ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, documentation that the patient is homozygous for the F508del mutation in the CFTR gene provided from an FDA-cleared CF mutation test, attestation baseline and follow-up ophthalmologic exams will be performed in pediatric patients starting on therapy.
Age Restrictions	2 years of age or older
Prescriber Restrictions	Restricted to pulmonology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# OSMOLEX (amantadine)

---

## Products Affected

- OSMOLEX ER ORAL TABLET ER 24 HOUR THERAPY PACK
- OSMOLEX ER ORAL TABLET EXTENDED RELEASE 24 HOUR 129 MG, 193 MG, 258 MG

PA Criteria	Criteria Details
Exclusion Criteria	End-stage renal disease (creatinine clearance below 15 mL/min)
Required Medical Information	Diagnosis of covered use, documentation patient tried and failed immediate-release amantadine.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to neurology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	



# OXBRYTA (voxelotor)

---

## Products Affected

- OXBRYTA

PA Criteria	Criteria Details
Exclusion Criteria	Hemoglobin greater than 10.5 g/dL
Required Medical Information	Diagnosis of covered use, submission of hemoglobin level.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to hematology
Coverage Duration	6 months
Other Criteria	PA applies to all. Submission of improved hemoglobin level from baseline will be required for initial reauthorization after the first 6 months. Documentation of continued hemoglobin level improvement or maintenance of initial hemoglobin level improvement will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# OXERVATE (cenegermin-bkbj)

---

## Products Affected

- OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	Restricted to optometry and ophthalmology
Coverage Duration	8 weeks
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# PANRETIN (alitretinoin)

---

## Products Affected

- PANRETIN

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, requirement for systemic Kaposi's sarcoma therapy (more than 10 new Kaposi's Sarcoma lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary Kaposi's sarcoma, or symptomatic visceral involvement)
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to dermatology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PARKINSON'S DISEASE "OFF" EPISODE (AS NEEDED) THERAPIES

## Products Affected

- APOKYN SUBCUTANEOUS SOLUTION CARTRIDGE
- INBRIJA
- KYNMOBI
- KYNMOBI TITRATION KIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For Inbrija, administration of non-selective monoamine oxidase inhibitors within 14 days of initiation, asthma, COPD, or other chronic underlying lung disease. For Kynmobi, severe renal impairment or end-stage renal disease, severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis of covered use, attestation patient is experiencing "off" episodes despite carbidopa/levodopa therapy, prescription claims or documentation from physician showing patient is (a) currently taking at least one other medication to help reduce "off" episodes (from COMT inhibitor, dopamine agonist, or monoamine oxidase B inhibitor drug classes) or (b) has tried and failed, had an intolerance to, or has a contraindication to medications from at least two different drug classes (COMT inhibitors, dopamine agonists, monoamine oxidase B inhibitors) that can help reduce "off" episodes.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PCSK9 INHIBITORS

## Products Affected

- PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- REPATHA
- REPATHA PUSHTRONEX SYSTEM
- REPATHA SURECLICK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	For all indications, submission of LDL level obtained within the previous 6 months. For primary hyperlipidemia (including HeFH) and ASCVD indications, submission of current or previous lipid-lowering therapies. For HeFH, documentation of genetic test result documenting HeFH or diagnosis by clinical criteria using Simon Broom or WHO/Dutch Lipid Network criteria. For ASCVD, documented history of MI, ACS, stable or unstable angina, coronary or other arterial revascularization, stroke, TIA, or PAD.
<b>Age Restrictions</b>	For Repatha, 13 years of age or older. For Praluent, 18 years of age or older.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. For approval for primary hyperlipidemia (including HeFH) and ASCVD indications, the patient must currently be using a statin plus ezetimibe or the patient must have tried and failed to have an adequate response to or had an intolerance to at least two statins or one statin and ezetimibe. At least one statin previously tried and failed must be a hydrophilic statin.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

## PDE5 INHIBITORS (PAH)

---

### Products Affected

- *alyq*
- *sildenafil citrate oral suspension reconstituted*
- *sildenafil citrate oral tablet 20 mg*
- *tadalafil (pah)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For tadalafil, diagnosis of severe (Child-Pugh class C) hepatic impairment, creatinine clearance below 30 mL/min or on hemodialysis
<b>Required Medical Information</b>	Diagnosis of covered use. For tadalafil, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to cardiology and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PEGFILGRASTIM

---

## Products Affected

- NEULASTA ONPRO
- NEULASTA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- UDENYCA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of FDA-approved indication.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	

# PEMAZYRE (pemigatinib)

---

## Products Affected

- PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A4 inducers
Required Medical Information	Diagnosis of covered use, submission of test result confirming presence of FGFR2 fusion or rearrangement, submission of previous systemic treatment(s) tried, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# PENICILLAMINE

---

## Products Affected

- *penicillamine oral capsule*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy unless being treated for Wilson's disease or certain patients with cystinuria, rheumatoid arthritis patients with a history or other evidence of renal insufficiency
<b>Required Medical Information</b>	Diagnosis of covered use, laboratory analysis applicable to indication for use, documentation that patient has tried and failed or had an intolerance to penicillamine tablets.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PIQRAY (alpelisib)

## Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing showing that the primary tumor type is HR-positive, HER2-negative, PIK3CA-mutated, attestation that patient has advanced or metastatic disease and will be taking concurrently with fulvestrant, submission of prior therapies tried, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# POMALYST (pomalidomide)

---

## Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, submission of baseline serum bilirubin, AST, ALT, CBC including ANC and platelet count, prior therapies, when prior therapy was completed, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# PRETOMANID

---

## Products Affected

- PRETOMANID

PA Criteria	Criteria Details
Exclusion Criteria	Inability to use bedaquiline or linezolid
Required Medical Information	Diagnosis of covered use, attestation pretomanid will be used in combination with bedaquiline and linezolid.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to infectious diseases and pulmonology.
Coverage Duration	26 weeks
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# PREVYMIS (letefmovir)

---

## Products Affected

- PREVYMIS ORAL

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, coadministration with ergot alkaloids, pimoziide, pitavastatin, or simvastatin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of day number post-HSCT, documentation of any previous doses of letermovir.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology, oncology, transplant specialist, and infectious diseases
<b>Coverage Duration</b>	100 days
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PRIOR AUTHORIZATION TO OVERRIDE SPECIALTY RESTRICTIONS

## Products Affected

- FABIOR
- PEG-INTRON REDIPEN SUBCUTANEOUS KIT 50 MCG/0.5ML
- PEG-INTRON SUBCUTANEOUS KIT 50 MCG/0.5ML
- PEGASYS PROCLICK SUBCUTANEOUS SOLUTION 135 MCG/0.5ML
- PEGASYS SUBCUTANEOUS SOLUTION
- PEGINTRON SUBCUTANEOUS KIT 50 MCG/0.5ML
- *tazarotene external*
- TAZORAC EXTERNAL CREAM 0.05 %
- TAZORAC EXTERNAL GEL
- VABOMERE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	The following physician specialties are exempt from prior authorization (by drug): (a) for Fabior, tazarotene, and Tazorac: dermatology exempt, (b) for Pegasys: gastroenterology, hepatology, or infectious diseases exempt, (c) for Vabomere: infectious diseases or nephrology exempt
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PROCYSBI (cysteamine)

---

## Products Affected

- PROCYSBI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation that patient has tried and failed or had an intolerance to immediate-release cysteamine.
Age Restrictions	1 year of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. For approval, the patient must have tried and failed to have an adequate response to, had an intolerance to, or have a contraindication to therapy with immediate-release cysteamine.
Indications	All Medically-accepted Indications.
Off Label Uses	

# PROLIA (denosumab)

## Products Affected

- PROLIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Hypocalcemia, pregnancy
<b>Required Medical Information</b>	Diagnosis of covered use, submission of calcium level, pregnancy status for female patients of childbearing potential. "High risk for fracture" is defined as (1) a history of osteoporotic fracture or (2) multiple risk factors for fracture or (3) patients who have failed or are intolerant of other available osteoporosis therapies, confirmation of osteoporosis diagnosis either through densitometry (T-score less than or equal to -2.5 at the total hip, femoral neck, or lumbar spine) or clinically (documented presence of fragility fracture).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated serum calcium level since the previous authorization will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# PROMACTA (eltrombopag)

---

## Products Affected

- PROMACTA ORAL PACKET
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use (including cause of thrombocytopenia if being used for that indication), documentation of previous therapies tried (corticosteroids, immunoglobulins), submission of platelet count.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	PA applies to all. Updated platelet count since the previous authorization will be required for subsequent reauthorizations. Not indicated for treatment of patients with myelodysplastic syndrome and will not be approved for this use.
Indications	All Medically-accepted Indications.
Off Label Uses	

# PROMETHAZINE IN OLDER PATIENTS

## Products Affected

- *promethazine hcl oral*
- PROMETHAZINE HCL RECTAL SUPPOSITORY 12.5 MG
- *promethazine hcl rectal suppository 25 mg*
- *promethazine vc plain*
- *promethazine-phenylephrine*
- *promethegan rectal suppository 25 mg, 50 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use. For allergic conditions, documentation must be submitted showing patient has tried and failed or had an inadequate response to a second-generation antihistamine.
<b>Age Restrictions</b>	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	Promethazine is a potent anticholinergic considered high-risk in older patients due to risks of confusion, dry mouth, constipation, and decreased clearance with advanced age.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# PROSTATE CANCER ORAL MEDICATIONS

## Products Affected

- *abiraterone acetate oral tablet 250 mg*
- ERLEADA
- NUBEQA
- XTANDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	For abiraterone, severe hepatic impairment (Child-Pugh class C)
<b>Required Medical Information</b>	Diagnosis of covered use. For Nubeqa, documentation of other treatments tried. For abiraterone, confirmation patient will receive concurrent prednisone, submission of baseline ALT, AST, bilirubin, and serum potassium level.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology and urology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. Nubeqa will be authorized only if the patient previously tried and had an inadequate clinical response or an intolerance to both Erleada and Xtandi.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# QINLOCK (ripretinib)

---

## Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of previous kinase inhibitor therapies, baseline blood pressure reading, attestation patient does not have uncontrolled hypertension, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# RAVICTI (glycerol phenylbutyrate)

---

## Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline fasting plasma ammonia level.
Age Restrictions	2 months of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# RETACRIT (epoetin alfa-epbx)

## Products Affected

- RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, submission of hemoglobin or hematocrit level, serum iron, total iron-binding capacity (TIBC), and transferrin within 30 days of request date, documentation that the patient does not have uncontrolled hypertension.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	For non-ESRD-related conditions: 90 days. For ESRD-related conditions: 1 year.
<b>Other Criteria</b>	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# RETEVMO (selpercatinib)

---

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncontrolled hypertension, coadministration with moderate or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of FDA-approved test confirming presence of RET gene fusion or mutation, baseline blood pressure reading, pregnancy status for female patients of childbearing potential. For patients with RET fusion-positive thyroid cancer, submission of date or year of previous previous radioactive iodine treatment or reason why radioactive iodine therapy is not appropriate.
<b>Age Restrictions</b>	12 years of age or older based on indication
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# REVCOVI (elapegademase-lvlr)

---

## Products Affected

- REVCOVI

PA Criteria	Criteria Details
Exclusion Criteria	Severe thrombocytopenia
Required Medical Information	Diagnosis of covered use, submission of plasma ADA activity and platelet count.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated plasma ADA level and platelet count since the previous authorization will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All FDA-approved Indications.
Off Label Uses	



# REVLIMID (lenalidomide)

---

## Products Affected

- REVLIMID

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, chronic lymphocytic leukemia (outside of a controlled clinical trial)
<b>Required Medical Information</b>	Diagnosis of covered use, submission of CBC including ANC and platelet count, pregnancy status for female patients of childbearing potential. For mantle cell lymphoma, documentation of at least two prior therapies tried, one of which included bortezomib (or a documented contraindication to bortezomib). For follicular lymphoma and marginal zone lymphoma, submission of prior treatments tried and attestation medication will be coadministered with a rituximab product.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# REZUROCK (belumosudil)

---

## Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of at least 2 previous therapies tried and failed for chronic graft-versus-host disease, pregnancy status for female patients of childbearing potential.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to hematology, oncology, and transplant specialty
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All FDA-approved Indications.
Off Label Uses	

# ROZLYTREK (entrectinib)

---

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential. For non-small cell lung cancer, submission of results showing tumor is ROS1-positive as detected by an FDA-approved test. For solid tumors, submission of evidence of a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# RUBRACA (rucaparib)

---

## Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline CBC. For BRCA mutation-associated ovarian, fallopian tube, primary peritoneal or metastatic castration-resistant prostate cancer, confirmation of deleterious BRCA mutation as detected by FDA-approved companion diagnostic test, documentation that the patient has been treated with two or more chemotherapies.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology and urology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only. For initial approval, patient must have recovered from hematological toxicity caused by previous chemotherapy (Grade 1 or less).
Indications	All Medically-accepted Indications.
Off Label Uses	

# RYDAPT (midostaurin)

---

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use. For acute myeloid leukemia, submission of FDA-approved test confirming presence of FLT3 mutation, documentation of other chemotherapy that will be coadministered with midostaurin.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SAPROPTERIN

---

## Products Affected

- *sapropterin dihydrochloride oral packet*
- *sapropterin dihydrochloride oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of blood phenylalanine level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to all. Submission of blood phenylalanine concentration documenting a reduction in blood phenylalanine concentration from pre-treatment baseline will be required for initial reauthorization. Documentation of continued phenylalanine level improvement or maintenance of initial phenylalanine level improvement will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SEDATIVE HYPNOTICS IN OLDER PATIENTS

---

## Products Affected

- *eszopiclone*
- *zaleplon*
- *zolpidem tartrate oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, documentation at least two of the following medications were tried and deemed ineffective or intolerable: Belsomra, doxepin tablets, ramelteon, and trazodone.
<b>Age Restrictions</b>	PA applies to patients 65 years of age or older. PA does not apply to patients 64 years of age or younger.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only. Sedative hypnotic medications are high-risk medications in older patients due to increased risks of cognitive impairment, delirium, unsteady gait, syncope, falls, fractures, and motor vehicle accidents.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SEROSTIM (somatropin)

---

## Products Affected

- SEROSTIM SUBCUTANEOUS SOLUTION  
RECONSTITUTED 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	Active malignancy, acute critical illness, active proliferative or severe non-proliferative diabetic retinopathy
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only. Serostim is indicated only for the treatment of HIV-associated cachexia/wasting and uses outside of this indication will not be approved.
Indications	All Medically-accepted Indications.
Off Label Uses	



# SIGNIFOR (pasireotide)

## Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment (Child-Pugh class C), uncorrected hypokalemia or hypomagnesemia
<b>Required Medical Information</b>	Diagnosis of covered use, submission of ALT, aspartate aminotransferase, alkaline phosphatase, total bilirubin, and serum potassium and magnesium levels, attestation that pituitary radiation/surgery was not curative or is not an option.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to endocrinology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated liver function testing and serum potassium and magnesium levels since the previous authorization will be required for subsequent reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SIMVASTATIN 80 mg per day

---

## Products Affected

- *ezetimibe-simvastatin oral tablet 10-80 mg*
- *simvastatin oral tablet 80 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Any new start to therapy. Not recommended as initial therapy nor for patients already taking lower doses of simvastatin whose response is inadequate.
<b>Required Medical Information</b>	Diagnosis of covered use, documentation that patient has been taking simvastatin 80 mg daily for 12 months or longer without ill effect, submission of lipid panel, liver function tests, and serum creatinine level all obtained within the past 12 months.
<b>Age Restrictions</b>	10 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SIRTURO (bedaquiline)

---

## Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline ECG, serum potassium, calcium, magnesium, ALT, AST, alkaline phosphatase, and bilirubin, confirmation that Sirturo will be co-administered with pretomanid and linezolid or at least 3 other drugs proven to be or at least 4 other drugs suspected to be effective against the patient's M. tuberculosis isolate and submission of susceptibility testing, if available.
Age Restrictions	5 years of age or older
Prescriber Restrictions	Restricted to infectious diseases and pulmonology
Coverage Duration	26 weeks
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SIVEXTRO (tedizolid)

---

## Products Affected

- SIVEXTRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of a culture and sensitivity showing that the suspected causative agent is susceptible to this medication.
Age Restrictions	12 years of age or older
Prescriber Restrictions	Restricted to infectious diseases
Coverage Duration	6 days
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SOMATULINE DEPOT (lanreotide)

---

## Products Affected

- SOMATULINE DEPOT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of baseline serum GH, IGF-1, TSH, and blood glucose levels. For acromegaly, attestation that pituitary radiation/surgery was not curative or is not an option.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology and oncology
Coverage Duration	Initially 3 months, then up to 1 year
Other Criteria	PA applies to new starts only. Continuation of therapy requires documentation of a positive clinical response. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SOMAVERT (pegvisomant)

---

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline IGF-1, ALT, AST, alkaline phosphatase, and serum total bilirubin, attestation that surgery or radiation was not curative or is not an option.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to endocrinology
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated IGF-1 level and liver function tests since the previous authorization will be required for subsequent annual reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SOVALDI (sofosbuvir)

---

## Products Affected

- SOVALDI ORAL PACKET
- SOVALDI ORAL TABLET 200 MG, 400 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) genotype 1a, 1b, 2, 3, or 4 infection, documentation of whether cirrhosis is present or not and whether or not it is compensated or decompensated, confirmation a test for HBV infection (HBsAg and anti-HBc) was completed.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Criteria for coverage duration will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SPRYCEL (dasatinib)

---

## Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	Uncorrected hypokalemia, uncorrected hypomagnesemia
Required Medical Information	Diagnosis of covered use, submission of serum potassium and magnesium, pregnancy status for female patients of childbearing potential. For adults with resistance or intolerance to prior therapy, documentation of prior therapy.
Age Restrictions	
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# STIVARGA (regorafenib)

---

## Products Affected

- STIVARGA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe or uncontrolled hypertension, coadministration with strong CYP3A4 inhibitors or inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of previous therapies, submission of baseline ALT, AST, serum bilirubin, blood pressure reading, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SUCRAID (sacrosidase)

---

## Products Affected

- SUCRAID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of laboratory-confirmed congenital sucrase-isomaltase deficiency via differential urinary disaccharide test or measurement of intestinal disaccharides following small bowel biopsy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SUTENT (sunitinib)

---

## Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SYMDEKO (tezacaftor/ivacaftor)

---

## Products Affected

- SYMDEKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use. Documentation that the patient is homozygous for the F508del mutation or has at least one mutation in the CFTR gene responsive to the drug (see section 12.1, table 4 of package insert for full list) provided from an FDA-cleared CF mutation test. Submission of documentation that baseline and follow-up ophthalmologic exams will be performed in pediatric patients starting on therapy.
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	Restricted to pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SYMLIN (pramlintide)

---

## Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	Gastroparesis
Required Medical Information	Diagnosis of covered use, confirmation of current use of a mealtime insulin.
Age Restrictions	
Prescriber Restrictions	PA not required for endocrinology
Coverage Duration	1 year
Other Criteria	PA applies to all except when prescribed by endocrinology.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SYMPROIC (naldemedine)

---

## Products Affected

- SYMPROIC

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Known or suspected gastrointestinal obstruction or increased risk of recurrent obstruction, severe hepatic impairment (Child-Pugh class C)
<b>Required Medical Information</b>	Diagnosis of covered use, documentation patient has been using opioids at a morphine equivalent dose of at least 30 mg daily for at least 4 weeks prior to initiation, provider must attest that if opioid medication is stopped for any reason, naldemedine will be discontinued.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# SYNAREL (nafarelin)

---

## Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy/breast-feeding, undiagnosed abnormal vaginal bleeding
Required Medical Information	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	For endometriosis, 6 months. For all other diagnoses, 1 year.
Other Criteria	PA applies to all. Re-treatment for endometriosis is not recommended because safety data are not available.
Indications	All Medically-accepted Indications.
Off Label Uses	

# SYNRIBO (omacetaxine)

---

## Products Affected

- SYNRIPO

PA Criteria	Criteria Details
Exclusion Criteria	Poor glycemic control
Required Medical Information	Diagnosis of covered use, submission of prior therapies tried and failed, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	



# TABRECTA (capmatinib)

---

## Products Affected

- TABRECTA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with moderate or strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TAFAMIDIS

---

## Products Affected

- VYNDAMAX
- VYNDAQEL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TAFINLAR (dabrafenib)

---

## Products Affected

- TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP2C8 or CYP3A4 inhibitors
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of BRAF V600E or V600K mutation. For non-small cell lung cancer, thyroid cancer, or unresectable/metastatic melanoma with a BRAF V600K mutation, attestation that therapy will be used in combination with trametinib.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TAGRISSO (osimertinib)

---

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, confirmation of the presence of required genetic mutations/deletions as detected by an FDA-approved test, pregnancy status for female patients of childbearing potential. For EGFR T790M mutation-positive NSCLC, documentation that the patient has progressed on or after EGFR TKI therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	Initially 3 months, then 1 year
Other Criteria	PA applies to new starts only. Continuation of approval requires affirmation of absence of unacceptable toxicities.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TALZENNA (talazoparib)

---

## Products Affected

- TALZENNA ORAL CAPSULE 0.25 MG, 1 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of FDA-approved companion test results showing patient is a candidate for therapy and pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TARGRETIN (bexarotene) GEL

---

## Products Affected

- TARGRETIN EXTERNAL

PA Criteria	Criteria Details
Exclusion Criteria	Pregnancy
Required Medical Information	Diagnosis of covered use, submission of previous therapies.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All FDA-approved Indications.
Off Label Uses	

# TASIGNA (nilotinib)

---

## Products Affected

- TASIGNA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia, uncorrected hypomagnesemia, long QT syndrome, coadministration with drugs that prolong the QT interval or strong CYP3A4 inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline ECG, Philadelphia chromosome (Ph) status, potassium and magnesium levels.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TAVALISSE (fostamatinib)

---

## Products Affected

- TAVALISSE ORAL TABLET 100 MG, 150 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of platelet count, documentation patient had an insufficient response to prior treatment (including at least one of the following: corticosteroids, immunoglobulins, splenectomy, and/or a thrombopoietin receptor agonist).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology
<b>Coverage Duration</b>	Initially 12 weeks, then 1 year
<b>Other Criteria</b>	PA applies to all. Documentation of an improvement in platelet count will be required for initial reauthorization after the first 12 weeks. Maintenance of a clinical benefit will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# TAZVERIK (tazemetostat)

---

## Products Affected

- TAZVERIK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inhibitors or moderate or strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	16 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	Initially 16 weeks, then 1 year
<b>Other Criteria</b>	PA applies to new starts only. Continuation of therapy requires (a) documentation of a positive clinical response and (b) attestation no known secondary malignancies have developed.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TEGSEDI (inotersen)

---

## Products Affected

- TEGSEDI

PA Criteria	Criteria Details
Exclusion Criteria	Platelet count less than $100 \times 10^9$ L
Required Medical Information	Diagnosis of covered use, submission of platelet count.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated platelet count since the previous authorization will be required for subsequent annual reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TEPMETKO (tepotinib)

---

## Products Affected

- TEPMETKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with strong CYP3A inducers or dual strong CYP3A4/P-glycoprotein inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of FDA-approved test confirming presence of MET exon 14 skipping mutation, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TESTOSTERONE REPLACEMENT PRODUCTS

## Products Affected

- ANDRODERM TRANSDERMAL PATCH 24 HOUR (1%)
- TESTOSTERONE TRANSDERMAL GEL 10 MG/ACT (2%) • *testosterone transdermal solution*
- *testosterone transdermal gel 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of serum testosterone level, documentation that patient has been evaluated for the presence of breast and prostate cancer prior to initiation of therapy.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Documentation of clinically relevant response to therapy (including, but not limited to submission of updated serum testosterone level) will be required for subsequent reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TIBSOVO (ivosidenib)

---

## Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of IDH1 mutation.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TOBI PODHALER (tobramycin)

---

## Products Affected

- TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	6 years of age or older
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TOLVAPTAN (HYPONATREMIA)

---

## Products Affected

- SAMSCA ORAL TABLET 15 MG
- *tolvaptan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of signs or symptoms of significant liver impairment or injury, need to raise serum sodium acutely, inability to sense or respond to thirst, hypovolemia, anuria, coadministration with strong CYP3A inhibitors or inducers or desmopressin
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum sodium.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	30 days
<b>Other Criteria</b>	PA applies to all. Treatment should be initiated in a setting where serum sodium can be monitored closely. Treatment is limited to 30 days to prevent liver injury. This formulation of tolvaptan will not be approved for autosomal dominant polycystic kidney disease (ADPKD) because the tolvaptan formulation approved for ADPKD has a mandatory REMS program.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TOPICAL ONYCHOMYCOSIS TREATMENTS

---

## Products Affected

- JUBLIA
- *tavaborole*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, submission of culture-proven <i>Trichophyton rubrum</i> or <i>Trichophyton mentagrophytes</i> infection, documentation patient has (1) tried and failed to respond to or tolerate oral terbinafine therapy or a documented contraindication to its use exists, and (2) tried and failed therapy with topical ciclopirox nail solution.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	48 weeks
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# TRIENTINE

---

## Products Affected

- *clovique*
- *trientine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of intolerance to penicillamine.
Age Restrictions	6 years of age or older
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# TRUSELTIQ (infigratinib)

---

## Products Affected

- TRUSELTIQ (100MG DAILY DOSE)
- TRUSELTIQ (125MG DAILY DOSE)
- TRUSELTIQ (50MG DAILY DOSE)
- TRUSELTIQ (75MG DAILY DOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inducers or inhibitors, coadministration with proton pump inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use, submission of test result confirming presence of FGFR2 fusion or rearrangement, submission of previous systemic treatment(s) tried.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TUKYSA (tucatinib)

---

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Pregnancy, coadministration with strong CYP3A inducers, strong CYP2C8 inhibitors, or moderate CYP2C8 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing showing that the primary tumor type is HER2-positive, submission of previous systemic treatment including prior HER2-directed therapy, pregnancy status for female patients of childbearing potential, confirmation that the treatment regimen will include concomitant use of capecitabine and trastuzumab.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TURALIO (pexidartinib)

---

## Products Affected

- TURALIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Active liver or biliary tract disease (including increased ALP), pre-existing increased serum transaminases, total or direct bilirubin greater than the upper limit of normal, coadministration with other hepatotoxic medications, strong CYP3A inducers, or proton pump inhibitors
<b>Required Medical Information</b>	Diagnosis of covered use (and surgical intervention is not possible or practical), submission of serum transaminases, total and direct bilirubin, and ALP.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# TYMLOS (abaloparatide)

## Products Affected

- TYMLOS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Male patients, female patients of childbearing potential, pre-existing hypercalcemia, underlying hypercalcemic disorder (such as primary hyperparathyroidism), patients with an increased risk of osteosarcoma (such as those with Paget's disease)
<b>Required Medical Information</b>	Diagnosis of covered use where "high risk for fracture" is defined as (1) a history of osteoporotic fracture or (2) multiple risk factors for fracture or (3) patients who have failed or are intolerant of other available osteoporosis therapies, submission of baseline serum calcium, postmenopausal status, documentation that at least one bisphosphonate was tried and failed (or a bisphosphonate is contraindicated), submission of a value, condition, or past medical history that assesses fracture risk (e.g., DEXA scan results or prior fracture), submission of number of total months of all prior use of parathyroid hormone analogs and parathyroid hormone related peptides.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	2 years maximum dependent on patient's prior use of all PTH analogs and PTH-related peptides
<b>Other Criteria</b>	PA applies to all. Use of parathyroid hormone analogs and/or parathyroid hormone related peptides for more than 2 years during a patient's lifetime is not recommended and requests for therapy with any of these agents for a combined total of greater than 2 years will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# UKONIQ (umbralisib)

---

## Products Affected

- UKONIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential, attestation patient will receive prophylaxis for <i>Pneumocystis jirovecii</i> pneumonia (PJP) and, if necessary, cytomegalovirus. For follicular lymphoma, submission of at least three prior systemic therapies used. For marginal zone lymphoma, submission of at least one prior anti-CD20-based regimen used.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

## UPNEEQ (oxymetazoline)

---

### Products Affected

- UPNEEQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to ophthalmologic surgery
Coverage Duration	Initially 90 days, then 1 year
Other Criteria	PA applies to all. Submission of clinically significant response to therapy will be required for reauthorization. Not FDA-approved for cosmetic use and therefore uses outside of acquired blepharoptosis will not be approved.
Indications	All FDA-approved Indications.
Off Label Uses	

# UPTRAVI (selexipag)

---

## Products Affected

- UPTRAVI ORAL TABLET 1000 MCG, 1200 MCG, 1400 MCG, 1600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	Severe (Child-Pugh class C) hepatic impairment
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	



# VALCHLOR (mechlorethamine)

---

## Products Affected

- VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	Use as initial therapy
Required Medical Information	Diagnosis of covered use, submission of previous skin-directed therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to dermatology and oncology
Coverage Duration	Initially 6 months, then 1 year
Other Criteria	PA applies to new starts only. Submission of clinically significant response to therapy will be required for reauthorization.
Indications	All Medically-accepted Indications.
Off Label Uses	

# VECAMEYL (mecamylamine)

---

## Products Affected

- VECAMEYL

PA Criteria	Criteria Details
Exclusion Criteria	Mild, moderate, labile hypertension, coronary insufficiency or history of recent myocardial infarction, uremia, glaucoma, organic pyloric stenosis
Required Medical Information	Diagnosis of covered use.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and neurology
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All Medically-accepted Indications.
Off Label Uses	

# VEMLIDY (tenofovir alafenamide)

---

## Products Affected

- VEMLIDY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	End-stage renal disease patients not receiving chronic hemodialysis, decompensated (Child-Pugh class B or C) hepatic impairment
<b>Required Medical Information</b>	Diagnosis of covered use, submission of creatinine clearance (or serum creatinine, actual body weight, and height to calculate estimated creatinine clearance), confirmation of HIV test and that drug will not be used by itself in the case of HIV co-infection.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	PA not required for gastroenterology or infectious diseases
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all except gastroenterology or infectious diseases. Updated creatinine clearance since the previous authorization will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VENCLEXTA (venetoclax)

---

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A inducers. For CLL/SLL, coadministration with strong CYP3A inhibitors at treatment initiation and initial dosage titration.
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VENTAVIS (iloprost)

---

## Products Affected

- VENTAVIS

PA Criteria	Criteria Details
Exclusion Criteria	Systolic blood pressure below 85 mmHg
Required Medical Information	Diagnosis of covered use, submission of baseline systolic blood pressure.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to cardiology and pulmonology
Coverage Duration	1 year
Other Criteria	This medication is covered as a Part B benefit except for enrollees residing in a long-term care facility. PA applies to new starts only when covered as a Part D benefit.
Indications	All Medically-accepted Indications.
Off Label Uses	

# VERQUVO (vericiguat)

## Products Affected

- VERQUVO ORAL TABLET 10 MG, 2.5 MG, 5 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Concomitant use of another soluble guanylate cyclase (sGC) stimulator or a phosphodiesterase-5 (PDE-5) inhibitor
<b>Required Medical Information</b>	Diagnosis, including either hospitalization for heart failure with reduced ejection fraction (HFrEF) within the previous 6 months or outpatient IV diuretic use within the previous 3 months, submission of left ventricular ejection fraction and pregnancy status for female patients of childbearing potential. Prescribers are also required to submit current regimen for the treatment of HFrEF, which must include (1) a renin-angiotensin system (RAS) inhibitor (ACE inhibitor, ARB, or sacubitril/valsartan), (2) a beta-blocker (BB), and (3) a mineralocorticoid receptor antagonist (MRA), each at maximally-tolerated doses. If any of these three therapies are not currently being used, prescriber is required to submit documentation as to why (e.g., contraindications, intolerances, etc.). Using the recommended dose of each therapeutic component to treat HFrEF is required. If the doses of any of these three components have not been optimized to the recommended dose to treat HFrEF, the prescriber is required to submit documentation as to why (e.g., intolerances, physiologic parameters, etc.). If the patient is using a BB not indicated for HFrEF, the patient will be required to switch to one of the three FDA-approved BBs for HFrEF (bisoprolol, carvedilol, or metoprolol succinate).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

# VERZENIO (abemaciclib)

---

## Products Affected

- VERZENIO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Coadministration with moderate or strong CYP3A4 inducers or ketoconazole
<b>Required Medical Information</b>	Diagnosis of covered use, submission of genetic tumor testing showing that the primary tumor type is HR-positive, HER2-negative, submission of baseline liver function tests and CBC, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VIBERZI (eluxadoline)

---

## Products Affected

- VIBERZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Prior cholecystectomy, known or suspected biliary duct obstruction, known or suspected sphincter of Oddi disease or dysfunction, alcoholism, alcohol abuse, alcohol addiction, or patients who drink more than 3 alcoholic beverages/day, history of pancreatitis, structural diseases of pancreas including known or suspected pancreatic duct obstruction, severe hepatic impairment (Child-Pugh class C), severe constipation or sequelae from constipation, known or suspected mechanical gastrointestinal obstruction
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to gastroenterology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	



# VITRAKVI (larotrectinib)

---

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of evidence the solid tumor has a neurotrophic receptor tyrosine kinase (NTRK) gene fusion without a known acquired resistance mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# VIZIMPRO (dacomitinib)

---

## Products Affected

- VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with a proton pump inhibitor
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of EGFR exon 19 deletion or exon 21 L858R substitution mutation, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# VMAT2 INHIBITORS

---

## Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG
- INGREZZA ORAL CAPSULE THERAPY PACK
- *tetrabenazine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Congenital long QT syndrome or a history of cardiac arrhythmia associated with a prolonged QT interval, coadministration with monoamine oxidase inhibitors. For tetrabenazine and Austedo, actively suicidal or untreated/undertreated depression, hepatic impairment. For Ingrezza, coadministration with strong CYP3A4 inducers.
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VOSEVI (sofosbuvir, velpatasvir, voxilaprevir)

---

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Moderate or severe hepatic impairment, coadministration with rifampin or drugs that are strong P-glycoprotein inducers or moderate to strong CYP2B6, CYP2C8, or CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use, laboratory confirmation of hepatitis C virus (HCV) and genotype, documentation of whether cirrhosis is present or not and whether or not it is compensated or decompensated, submission of previous treatment regimen, confirmation a test for HBV infection (HBsAg and anti-HBc) was completed.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 weeks
<b>Other Criteria</b>	PA applies to all.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VOTRIENT (pazopanib)

---

## Products Affected

- VOTRIENT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, uncontrolled hypertension, uncorrected hypokalemia, hypocalcemia, or hypomagnesemia, coadministration with strong CYP3A4 inducers or drugs that can prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of baseline blood pressure, ALT, AST, bilirubin, and serum potassium, calcium, and magnesium, pregnancy status for female patients of childbearing potential. For soft tissue sarcoma, submission of previous chemotherapy regimen(s).
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# VRAYLAR (cariprazine)

---

## Products Affected

- VRAYLAR ORAL CAPSULE
- VRAYLAR ORAL CAPSULE THERAPY PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Dementia-related psychosis, severe hepatic impairment, severe renal impairment (creatinine clearance less than 30 mL/min), coadministration with CYP3A4 inducers
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and psychiatry
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# WAKIX (pitolisant)

## Products Affected

- WAKIX ORAL TABLET 17.8 MG, 4.45 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Severe hepatic impairment, end-stage renal disease, known QT interval prolongation, symptomatic bradycardia, uncorrected hypokalemia or hypomagnesemia, coadministration with medications that prolong the QT interval
<b>Required Medical Information</b>	Diagnosis of covered use, submission of serum potassium and magnesium.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology and sleep medicine
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated serum potassium and magnesium since the previous authorization will be required for subsequent annual reauthorizations. For excessive daytime sleepiness associated with narcolepsy, pitolisant will be authorized only if the patient previously tried and had an inadequate clinical response or an intolerance to armodafinil or modafinil.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# WHITE BLOOD CELL STIMULATORS

---

## Products Affected

- NIVESTYM
- ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	For approval of Nivestym, the patient must have tried and failed to have an adequate response to or had an intolerance to Zarxio. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	



# XALKORI (crizotinib)

---

## Products Affected

- XALKORI

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome, coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of results showing tumor is ALK or ROS1-positive as detected by an FDA-approved test, pregnancy status for female patients of childbearing potential.
Age Restrictions	For ALK-positive systemic anaplastic large cell lymphoma only, 1 year of age to 21 years of age
Prescriber Restrictions	Restricted to hematology and oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# XCOPRI (cenobamate)

---

## Products Affected

- XCOPRI (250 MG DAILY DOSE)
- XCOPRI (350 MG DAILY DOSE)
- XCOPRI ORAL TABLET 100 MG, 150 MG, 200 MG, 50 MG
- XCOPRI ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Familial short QT syndrome, patients with end-stage renal disease (creatinine clearance less than 15 mL/min) undergoing dialysis, severe hepatic impairment
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to neurology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# XERMELO (telotristat)

---

## Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation patient has been on at least 12 weeks of prior somatostatin analog therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	12 weeks
Other Criteria	PA applies to all. Continuation of therapy requires that symptoms have stabilized or improved and that the patient has not experienced episodes of severe constipation.
Indications	All Medically-accepted Indications.
Off Label Uses	

# XGEVA (denosumab)

---

## Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia
Required Medical Information	Diagnosis of covered use, submission of serum calcium level, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to new starts only. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# XOLAIR (omalizumab)

## Products Affected

- XOLAIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Patients whose pre-treatment serum IgE level and body weight place them in the "insufficient data to recommend a dose" category based on dosing charts in the prescribing information
<b>Required Medical Information</b>	Diagnosis of covered use. For asthma, documentation that patient's symptoms are poorly controlled with inhaled corticosteroids, submission of patient's current body weight, pre-treatment serum IgE level, pulmonary function test results including FEV1, positive skin test result or demonstrated in vitro reactivity (RAST test) to a perennial aeroallergen, frequency of inhaled short-acting beta2-agonist therapy, frequency of daily and nighttime symptoms and exacerbations, and effect of exacerbations on activity. For chronic idiopathic urticaria, documentation that the patient continues to experience severe itching and hives despite the use of an H1 antihistamine at an approved dose. For nasal polyps, documentation that patient's symptoms are poorly controlled with intranasal corticosteroids and current intranasal corticosteroid therapy.
<b>Age Restrictions</b>	6 years of age or older
<b>Prescriber Restrictions</b>	Restricted to allergy, dermatology, immunology, otolaryngology/otorhinolaryngology, and pulmonology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Submission of objective documentation of symptomatic improvement and updated patient weight will be required for subsequent annual reauthorizations. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# XOSPATA (gilteritinib)

---

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Uncorrected hypokalemia or hypomagnesemia, coadministration with dual strong CYP3A/P-glycoprotein inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of FDA-approved test confirming presence of FLT3 mutation, serum calcium and magnesium, pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# XPOVIO (selinexor)

## Products Affected

- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG
- XPOVIO (80 MG TWICE WEEKLY)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	Diagnosis of covered use, documentation of treatment failure with or intolerance to all prior therapies to match the indication, submission of pregnancy status for female patients of childbearing potential.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

# XURIDEN (uridine triacetate)

---

## Products Affected

- XURIDEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, submission of baseline CBC including neutrophil count and mean corpuscular volume, baseline urine orotic acid level.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all. Updated urine orotic acid level and CBC including neutrophil count and mean corpuscular volume since the previous authorization will be required for subsequent annual reauthorizations.
Indications	All Medically-accepted Indications.
Off Label Uses	



# ZEJULA (niraparib)

---

## Products Affected

- ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use, documentation of treatment failure with or intolerance to all prior therapies to match the indication, submission of pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ZERBAXA (ceftolozane/tazobactam)

## Products Affected

- ZERBAXA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of covered use. For complicated intra-abdominal infections, confirmation patient will receive concurrent metronidazole therapy.
Age Restrictions	18 years of age or older
Prescriber Restrictions	
Coverage Duration	For UTI including pyelonephritis, 7 days. For all other FDA-approved indications, 14 days.
Other Criteria	PA applies to all. A description of how the drug will be used and/or obtained may be required to determine whether it will be covered under Medicare Part B or Part D. If the medication will be self-administered after proper training or the provider agrees to have the medication filled at a pharmacy and delivered to the office by the patient for provider administration, it may be covered under Part D. If these conditions are not satisfied this medication may be covered under Part B.
Indications	All Medically-accepted Indications.
Off Label Uses	

# ZILEUTON ER

---

## Products Affected

- ZILEUTON ER

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Active liver disease or persistent hepatic function elevation enzyme greater than or equal to 3 times the upper limit of normal
<b>Required Medical Information</b>	Diagnosis of covered use, submission of hepatic function enzymes and serum bilirubin.
<b>Age Restrictions</b>	12 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to all. Updated hepatic function enzymes and serum bilirubin since the previous authorization will be required for subsequent annual reauthorizations.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ZOKINVY (lonafarnib)

---

## Products Affected

- ZOKINVY ORAL CAPSULE 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	Body surface area less than 0.39 m <sup>2</sup>
Required Medical Information	Diagnosis of covered use including results of genetic testing supporting diagnosis, pregnancy status for female patients of childbearing potential.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 year
Other Criteria	PA applies to all.
Indications	All FDA-approved Indications.
Off Label Uses	

# ZONTIVITY (vorapaxar)

---

## Products Affected

- ZONTIVITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of stroke, transient ischemic attack, or intracranial hemorrhage, active pathological bleeding, severe hepatic impairment, coadministration with strong CYP3A inhibitors or inducers
<b>Required Medical Information</b>	Diagnosis of covered use, confirmation that patient has not had prior stroke, transient ischemic attack, or intracranial hemorrhage, documentation of concurrent use with aspirin and/or clopidogrel.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ZORBTIVE (somatropin)

---

## Products Affected

- ZORBTIVE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Active malignancy, acute critical illness, active proliferative or severe non-proliferative diabetic retinopathy
<b>Required Medical Information</b>	Diagnosis of covered use.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	4 weeks
<b>Other Criteria</b>	PA applies to all. Zorbtive is indicated only for the treatment of short bowel syndrome and uses outside of this indication will not be approved.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ZYDELIG (idelalisib)

---

## Products Affected

- ZYDELIG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	History of serious hypersensitivity reactions, including toxic epidermal necrolysis with any drug, coadministration with strong CYP3A inducers
<b>Required Medical Information</b>	Diagnosis of covered use, submission of pregnancy status for female patients of childbearing potential. For relapsed small lymphocytic lymphoma and follicular B-cell non-Hodgkin lymphoma, documentation of at least two prior systemic therapies.
<b>Age Restrictions</b>	18 years of age or older
<b>Prescriber Restrictions</b>	Restricted to hematology and oncology
<b>Coverage Duration</b>	1 year
<b>Other Criteria</b>	PA applies to new starts only.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	

# ZYKADIA (ceritinib)

---

## Products Affected

- ZYKADIA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	Coadministration with strong CYP3A inducers
Required Medical Information	Diagnosis of covered use, submission of FDA-approved test confirming presence of ALK-positive tumor, pregnancy status for female patients of childbearing potential.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Restricted to oncology
Coverage Duration	1 year
Other Criteria	PA applies to new starts only.
Indications	All Medically-accepted Indications.
Off Label Uses	





## Index

ABILIFY MYCITE MAINTENANCE KIT ORAL TABLET 10 MG, 15 MG, 20 MG, 30 MG, 5 MG.....	2	BUTALBITAL-APAP-CAFF-COD.....	26
ABILIFY MYCITE ORAL TABLET 10 MG, 15 MG, 20 MG, 30 MG, 5 MG.....	2	<i>butalbital-apap-caffeine oral capsule</i> .....	26
ABILIFY MYCITE STARTER KIT ORAL TABLET 10 MG, 15 MG, 20 MG, 30 MG, 5 MG.....	2	<i>butalbital-apap-caffeine oral tablet 50-325-40 mg</i> ....	26
<i>abiraterone acetate oral tablet 250 mg</i> .....	182	<i>butalbital-asa-caff-codeine</i> .....	26
ACTEMRA ACTPEN.....	20	<i>butalbital-aspirin-caffeine oral capsule</i> .....	26
ACTEMRA SUBCUTANEOUS.....	20	BUTISOL SODIUM ORAL TABLET 30 MG.....	89
ACTHAR.....	3	BYLVAY.....	27
ACTIMMUNE.....	4	BYLVAY (PELLETS).....	27
ADEMPAS.....	5	CABLIVI.....	28
AFINITOR DISPERZ.....	71	CABOMETYX.....	29
AFINITOR ORAL TABLET 10 MG.....	71	CALQUENCE.....	30
		CAPLYTA.....	31
		CAPRELSA.....	32
		CARBAGLU.....	33
		<i>carbinoxamine maleate oral solution</i> .....	77
		<i>carbinoxamine maleate oral tablet 4 mg</i> .....	77
		CARIMUNE NF INTRAVENOUS SOLUTION RECONSTITUTED 12 GM, 6 GM.....	96
AKYNZEO ORAL.....	6	CERDELGA.....	34
ALECENSA.....	7	CHENODAL.....	36
ALUNBRIG.....	9	CHLORPROPAMIDE ORAL TABLET 100 MG.....	89
<i>alyq</i> .....	169	CHOLBAM.....	37
<i>ambrisentan</i> .....	65	CIMZIA PREFILLED.....	20
ANADROL-50.....	11	CIMZIA STARTER KIT.....	20
ANDRODERM TRANSDERMAL PATCH 24 HOUR.....	226	CIMZIA SUBCUTANEOUS KIT 2 X 200 MG.....	20
APOKYN SUBCUTANEOUS SOLUTION CARTRIDGE....	167	<i>clemastine fumarate oral tablet 2.68 mg</i> .....	77
ARALAST NP INTRAVENOUS SOLUTION RECONSTITUTED 1000 MG, 500 MG.....	8	<i>clovique</i> .....	231
ARANESP (ALBUMIN FREE) INJECTION SOLUTION 100 MCG/ML, 200 MCG/ML, 25 MCG/ML, 300 MCG/ML, 40 MCG/ML, 60 MCG/ML.....	12	COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG.....	38
ARANESP (ALBUMIN FREE) INJECTION SOLUTION PREFILLED SYRINGE.....	12	COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG.....	38
ARCALYST.....	13	COMETRIQ (60 MG DAILY DOSE).....	38
ARIKAYCE.....	14	COPIKTRA ORAL CAPSULE 15 MG, 25 MG.....	39
<i>ascomp-codeine</i> .....	26	CORLANOR.....	40
AURYXIA.....	15		
AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG.....	250	<i>cyproheptadine hcl oral</i> .....	77
		CYSTADROPS.....	42
		CYSTARAN.....	42
		<i>dalfampridine er</i> .....	43
BAFIERTAM.....	79	DAURISMO ORAL TABLET 100 MG, 25 MG.....	44
BALVERSA.....	17	DAYVIGO ORAL TABLET 10 MG, 5 MG.....	45
BENLYSTA SUBCUTANEOUS.....	19	<i>deferasirox granules</i> .....	46
BIVIGAM INTRAVENOUS SOLUTION 5 GM/50ML.....	96	<i>deferasirox oral tablet</i> .....	46
<i>bosentan</i> .....	65	<i>deferasirox oral tablet soluble</i> .....	46
BOSULIF.....	21	<i>deferiprone</i> .....	47
BRAFTOVI ORAL CAPSULE 75 MG.....	22	DIACOMIT.....	48
		<i>diclofenac epolamine external</i> .....	52
BRONCHITOL.....	24	<i>diclofenac sodium external gel</i> .....	49, 51
BRUKINSA.....	25	<i>diclofenac sodium external solution</i> .....	50
BUPAP ORAL TABLET 50-300 MG.....	26	DIFICID.....	53
BUTALBITAL-ACETAMINOPHEN ORAL TABLET 50-300 MG, 50-325 MG.....	26	<i>digitek oral tablet 250 mcg</i> .....	54
		<i>digox oral tablet 250 mcg</i> .....	54

<i>digoxin oral tablet 250 mcg</i> .....	54	GATTEX .....	81
<i>diphenhydramine hcl oral elixir</i> .....	77	GAVRETO .....	82
<i>dronabinol</i> .....	56	GENOTROPIN .....	85
<i>droxidopa</i> .....	57	GENOTROPIN MINIQUICK .....	85
DUOBRII .....	58	GILOTRIF .....	83
DUOPA ENTERAL .....	59	GLASSIA .....	8
DUPIXENT .....	60	GOCOVRI .....	10
EGRIFTA SV .....	61	HAEGARDA .....	87
ELIGARD .....	84		
EMFLAZA .....	62	HETLIOZ .....	88
		HUMATROPE .....	85
EMPAVELI .....	63	<i>hydroxyzine hcl oral tablet</i> .....	77
EMSAM .....	64	<i>hydroxyzine pamoate oral</i> .....	77
ENSPRYNG .....	66	IBRANCE .....	90
EPCLUSA ORAL TABLET 200-50 MG, 400-100 MG ....	200	<i>icatibant acetate</i> .....	86
EPIDIOLEX .....	67	ICLUSIG .....	91
ERIVEDGE .....	68	IDHIFA .....	92
ERLEADA .....	182	ILARIS SUBCUTANEOUS SOLUTION .....	93
<i>erlotinib hcl</i> .....	69	IMBRUVICA .....	94
ESBRIET .....	70	IMCIVREE .....	95
<i>eszopiclone</i> .....	194	INBRIJA .....	167
<i>everolimus oral tablet 2.5 mg, 5 mg, 7.5 mg</i> .....	71	INCRELEX .....	132
		INGREZZA ORAL CAPSULE 40 MG, 60 MG, 80 MG ...	250
<i>ezetimibe-simvastatin oral tablet 10-80 mg</i> .....	197	INGREZZA ORAL CAPSULE THERAPY PACK .....	250
FABIOR .....	177	INLYTA .....	98
FARYDAK .....	73	INQOVI .....	99
		INREBIC .....	100
<i>fentanyl citrate buccal</i> .....	74	INTRON A .....	103
FERRIPROX ORAL SOLUTION .....	47	INVEGA HAFYERA .....	104
FERRIPROX ORAL TABLET 1000 MG .....	47	INVEGA TRINZA INTRAMUSCULAR SUSPENSION	
FINTEPLA .....	75	PREFILLED SYRINGE .....	105
FIRDAPSE .....	76	IRESSA .....	106
FIRMAGON (240 MG DOSE) .....	84	ISTURISA .....	107
FIRMAGON SUBCUTANEOUS SOLUTION		<i>itraconazole oral</i> .....	108
RECONSTITUTED 80 MG .....	84	JAKAFI .....	109
FLEBOGAMMA DIF .....	96	JUBLIA .....	230
		JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5	
FOTIVDA .....	78	MG .....	110
GALAFOLD .....	80	JYNARQUE .....	111
GAMMAGARD .....	96	KALYDECO .....	112
GAMMAGARD S/D LESS IGA .....	96	KERENDIA .....	113
GAMMAKED INJECTION SOLUTION 1 GM/10ML, 10		<i>ketoconazole oral</i> .....	114
GM/100ML, 20 GM/200ML, 5 GM/50ML .....	96	KEVEYIS .....	115
GAMMAPLEX INTRAVENOUS SOLUTION 10		KEVZARA .....	20
GM/100ML, 10 GM/200ML, 20 GM/200ML, 20			
GM/400ML, 5 GM/100ML, 5 GM/50ML .....	96	KORLYM .....	117
GAMUNEX-C .....	96	KOSELUGO .....	118

KYNMOBI .....	167	NAYZILAM .....	102
KYNMOBI TITRATION KIT .....	167	NERLYNX .....	146
<i>lapatinib ditosylate</i> .....	119	NEULASTA ONPRO .....	170
LENVIMA (10 MG DAILY DOSE) .....	121	NEULASTA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	170
LENVIMA (12 MG DAILY DOSE) .....	121	NEXAVAR .....	147
LENVIMA (14 MG DAILY DOSE) .....	121	NEXLETOL .....	18
LENVIMA (18 MG DAILY DOSE) .....	121	NEXLIZET .....	18
LENVIMA (20 MG DAILY DOSE) .....	121	NINLARO .....	148
LENVIMA (24 MG DAILY DOSE) .....	121	<i>nitisinone</i> .....	149
LENVIMA (4 MG DAILY DOSE) .....	121	NITYR .....	149
LENVIMA (8 MG DAILY DOSE) .....	121	NIVESTYM .....	255
LEUKINE INJECTION SOLUTION RECONSTITUTED .....	122	NORDITROPIN FLEXPRO SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	85
<i>leuprolide acetate injection</i> .....	84	NUBEQA .....	182
<i>lidocaine external patch 5 %</i> .....	123	NUCALA .....	101
LONSURF .....	124	NUDEXTA .....	150
LORBRENA ORAL TABLET 100 MG, 25 MG .....	125	NUPLAZID ORAL CAPSULE .....	151
LUCEMYRA .....	126	NUPLAZID ORAL TABLET 10 MG .....	151
LUMAKRAS .....	127	NUTROPIN AQ NUSPIN 10 SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	85
LUPANETA PACK .....	84	NUTROPIN AQ NUSPIN 20 SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	85
LUPKYNIS .....	128	NUTROPIN AQ NUSPIN 5 SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	85
LUPRON DEPOT (1-MONTH) .....	84	OCALIVA ORAL TABLET 10 MG, 5 MG .....	152
LUPRON DEPOT (3-MONTH) .....	84	OCTAGAM INTRAVENOUS SOLUTION 1 GM/20ML, 10 GM/100ML, 10 GM/200ML, 2 GM/20ML, 2.5 GM/50ML, 20 GM/200ML, 25 GM/500ML, 5 GM/100ML, 5 GM/50ML .....	96
LUPRON DEPOT (4-MONTH) .....	84	ODOMZO .....	153
LUPRON DEPOT (6-MONTH) .....	84	OFEV .....	154
LUPRON DEPOT-PED (1-MONTH) INTRAMUSCULAR KIT 11.25 MG, 15 MG .....	84	ONUREG .....	155
LUPRON DEPOT-PED (3-MONTH) INTRAMUSCULAR KIT 30 MG (PED) .....	84	OPSUMIT .....	65
LYNPARZA ORAL TABLET .....	129	ORENITRAM .....	156
MAVENCLAD (10 TABS) .....	130	ORFADIN ORAL CAPSULE 20 MG .....	149
MAVENCLAD (4 TABS) .....	130	ORFADIN ORAL SUSPENSION .....	149
MAVENCLAD (5 TABS) .....	130	ORGOVYX .....	157
MAVENCLAD (6 TABS) .....	130	ORIAHNN .....	158
MAVENCLAD (7 TABS) .....	130	ORLISSA ORAL TABLET 150 MG, 200 MG .....	159
MAVENCLAD (8 TABS) .....	130	ORKAMBI .....	160
MAVENCLAD (9 TABS) .....	130	ORLADEYO .....	87
MAVYRET .....	131	OSMOLEX ER ORAL TABLET ER 24 HOUR THERAPY PACK .....	161
<i>megestrol acetate oral suspension 40 mg/ml, 400 mg/10ml, 625 mg/5ml</i> .....	133	OSMOLEX ER ORAL TABLET EXTENDED RELEASE 24 HOUR 129 MG, 193 MG, 258 MG .....	161
MEKINIST .....	134	OTEZLA .....	20
MEKTOVI .....	22	OTREXUP SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.4ML, 12.5 MG/0.4ML, 15 MG/0.4ML, 17.5 MG/0.4ML, 20 MG/0.4ML, 22.5 MG/0.4ML, 25 MG/0.4ML .....	136
<i>meprobamate</i> .....	89	OXBRYTA .....	162
<i>methamphetamine hcl</i> .....	135		
METHITEST .....	138		
<i>methyltestosterone oral</i> .....	138		
<i>miglustat</i> .....	139		
MYALEPT .....	140		
MYCAPSSA .....	141		
MYFEMBREE .....	142		
MYTESI .....	143		
NAMZARIC .....	144		
NATPARA .....	145		

OXERVATE .....	163	REZUROCK .....	189
PANRETIN .....	166	ROZLYTREK ORAL CAPSULE 100 MG, 200 MG .....	190
PEGASYS PROCLICK SUBCUTANEOUS SOLUTION 135 MCG/0.5ML .....	177	RUBRACA .....	191
PEGASYS SUBCUTANEOUS SOLUTION .....	177	RUCONEST .....	86
PEG-INTRON REDIPEN SUBCUTANEOUS KIT 50 MCG/0.5ML .....	177	RYDAPT .....	192
PEGINTRON SUBCUTANEOUS KIT 50 MCG/0.5ML ...	177	SAMSCA ORAL TABLET 15 MG .....	229
PEG-INTRON SUBCUTANEOUS KIT 50 MCG/0.5ML ...	177	<i>sapropterin dihydrochloride oral packet</i> .....	193
PEMAZYRE ORAL TABLET 13.5 MG, 4.5 MG, 9 MG ...	171	<i>sapropterin dihydrochloride oral tablet</i> .....	193
<i>penicillamine oral capsule</i> .....	172	SEROSTIM SUBCUTANEOUS SOLUTION RECONSTITUTED 4 MG, 5 MG, 6 MG .....	195
PIQRAY (200 MG DAILY DOSE) .....	173	SIGNIFOR .....	196
PIQRAY (250 MG DAILY DOSE) .....	173	<i>sildenafil citrate oral suspension reconstituted</i> .....	169
PIQRAY (300 MG DAILY DOSE) .....	173	<i>sildenafil citrate oral tablet 20 mg</i> .....	169
POMALYST .....	174	SIMPONI SUBCUTANEOUS SOLUTION AUTO-INJECTOR .....	20
PRALUENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR .....	168	SIMPONI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	20
PRETOMANID .....	175	<i>simvastatin oral tablet 80 mg</i> .....	197
PREVYMIS ORAL .....	176	SIRTURO .....	198
PRIVIGEN .....	96	SIVEXTRO .....	199
PROCYSBI .....	178	SOMATULINE DEPOT .....	201
PROLASTIN-C INTRAVENOUS SOLUTION RECONSTITUTED .....	8	SOMAVERT .....	202
PROLIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .....	179	SOVALDI ORAL PACKET .....	203
PROMACTA ORAL PACKET .....	180	SOVALDI ORAL TABLET 200 MG, 400 MG .....	203
PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG .....	180	SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG .....	204
<i>promethazine hcl oral</i> .....	181	STIVARGA .....	205
PROMETHAZINE HCL RECTAL SUPPOSITORY 12.5 MG .....	181	SUCRAID .....	206
<i>promethazine hcl rectal suppository 25 mg</i> .....	181	<i>sunitinib malate</i> .....	208
<i>promethazine vc plain</i> .....	181	SYMDEKO .....	209
<i>promethazine-phenylephrine</i> .....	181	SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	210
<i>promethegan rectal suppository 25 mg, 50 mg</i> .....	181	SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR .....	210
QINLOCK .....	183	SYMPROIC .....	211
RASUVO SUBCUTANEOUS SOLUTION AUTO-INJECTOR 10 MG/0.2ML, 12.5 MG/0.25ML, 15 MG/0.3ML, 17.5 MG/0.35ML, 20 MG/0.4ML, 22.5 MG/0.45ML, 25 MG/0.5ML, 30 MG/0.6ML, 7.5 MG/0.15ML .....	136	SYNAREL .....	212
RAVICTI .....	184	SYNDROS .....	56
REDITREX .....	136	SYNRIBO .....	213
REPATHA .....	168	TABRECTA ORAL TABLET 150 MG, 200 MG .....	214
REPATHA PUSHTRONEX SYSTEM .....	168	<i>tadalafil (pah)</i> .....	169
REPATHA SURECLICK .....	168	TAFINLAR .....	216
RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML .....	185	TAGRISSE .....	217
RETEVMO ORAL CAPSULE 40 MG, 80 MG .....	186	<i>takhzyro</i> .....	87
REVCOVI .....	187	TALZENNA ORAL CAPSULE 0.25 MG, 1 MG .....	218
REVLIMID .....	188	TARGETIN EXTERNAL .....	219
		TASIGNA .....	220
		<i>tavorole</i> .....	230
		TAVALISSE ORAL TABLET 100 MG, 150 MG .....	221
		<i>tazarotene external</i> .....	177
		TAZORAC EXTERNAL CREAM 0.05 % .....	177
		TAZORAC EXTERNAL GEL .....	177

TAZVERIK .....	222	VERQUVO ORAL TABLET 10 MG, 2.5 MG, 5 MG .....	245
TEGSEDI .....	223	VERZENIO .....	246
TENCON ORAL TABLET 50-325 MG .....	26	VIBERZI .....	247
TEPMETKO .....	224	VITRAKVI ORAL CAPSULE 100 MG, 25 MG .....	248
<i>testosterone cypionate injection solution 200 mg/ml</i> .....	97	VITRAKVI ORAL SOLUTION .....	248
<i>testosterone cypionate intramuscular solution 100 mg/ml, 200 mg/ml</i> .....	97	VIZIMPRO ORAL TABLET 15 MG, 30 MG, 45 MG .....	249
<i>testosterone enanthate intramuscular solution</i> .....	97	VOSEVI .....	251
TESTOSTERONE TRANSDERMAL GEL 10 MG/ACT (2%) .....	226	VOTRIENT .....	252
<i>testosterone transdermal gel 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)</i> .....	226	VRAYLAR ORAL CAPSULE .....	253
<i>testosterone transdermal solution</i> .....	226	VRAYLAR ORAL CAPSULE THERAPY PACK .....	253
tetrabenazine .....	250	VTOL LQ .....	26
TIBSOVO .....	227	VUMERITY .....	79
TOBI PODHALER .....	228	VYNDAMAX .....	215
TOLSURA .....	108	VYNDAQEL .....	215
<i>tolvaptan</i> .....	229	WAKIX ORAL TABLET 17.8 MG, 4.45 MG .....	254
TRACLEER ORAL TABLET SOLUBLE .....	65	XALKORI .....	256
TRELSTAR MIXJECT .....	84	XCOPRI (250 MG DAILY DOSE) .....	258
TREMFYA .....	20	XCOPRI (350 MG DAILY DOSE) .....	258
<i>trientine hcl</i> .....	231	XCOPRI ORAL TABLET 100 MG, 150 MG, 200 MG, 50 MG .....	258
TRUSELTIQ (100MG DAILY DOSE) .....	233	XCOPRI ORAL TABLET THERAPY PACK .....	258
TRUSELTIQ (125MG DAILY DOSE) .....	233	XERMELO .....	259
TRUSELTIQ (50MG DAILY DOSE) .....	233	XGEVA .....	260
TRUSELTIQ (75MG DAILY DOSE) .....	233	XOLAIR .....	261
TUKYSA ORAL TABLET 150 MG, 50 MG .....	234	XOSPATA .....	262
TURALIO .....	235	XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 50 MG .....	263
TYMLOS .....	236	XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG .....	263
UDENYCA .....	170	XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG .....	263
UKONIQ .....	237	XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 60 MG .....	263
UPNEEQ .....	238	XPOVIO (60 MG TWICE WEEKLY) .....	263
UPTRAVI ORAL TABLET 1000 MCG, 1200 MCG, 1400 MCG, 1600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG .....	239	XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 20 MG, 40 MG .....	263
UPTRAVI ORAL TABLET THERAPY PACK .....	239	XPOVIO (80 MG TWICE WEEKLY) .....	263
VABOMERE .....	177	XTANDI .....	182
VALCHLOR .....	240	XURIDEN .....	264
VALTOCO 10 MG DOSE .....	102	<i>zaleplon</i> .....	194
VALTOCO 15 MG DOSE .....	102	ZARXIO .....	255
VALTOCO 20 MG DOSE .....	102	ZEBUTAL ORAL CAPSULE 50-325-40 MG .....	26
VALTOCO 5 MG DOSE .....	102	ZEJULA .....	265
VECAMYL .....	241	ZEMAIRA .....	8
VEMLIDY .....	242	ZEPOSIA .....	20
VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG ..	243	ZEPOSIA 7-DAY STARTER PACK .....	20
VENCLEXTA STARTING PACK .....	243	ZEPOSIA STARTER KIT .....	20
VENTAVIS .....	244	ZERBAXA .....	266
		ZILEUTON ER .....	267

ZOKINVY ORAL CAPSULE 50 MG, 75 MG .....	268
<i>zolpidem tartrate oral</i> .....	194
ZONTIVITY .....	269
ZORBIVE .....	270
ZYDELIG .....	271
ZYKADIA ORAL TABLET .....	272